

**Baseline Conditions Report  
for  
Ramona Grasslands Preserve  
San Diego County**



**January 2007**



"Funding for this project has been provided in full or in part through an Agreement with the State Water Resources Control Board (SWRCB) pursuant to the Costa-Machado Water Act of 2000 (Proposition 13) and any amendments thereto for the implementation of California's Non-point Source Pollution Control Program. The contents of this document do not necessarily reflect the views and policies of the SWRCB, nor does mention of trade names or commercial products constitute endorsement or recommendation for use."

**Baseline Conditions Report  
for  
Ramona Grasslands Preserve  
San Diego County**

*Prepared for:*

**County of San Diego**

Department of Parks and Recreation

9150 Chesapeake Drive, Suite 200

San Diego, CA 92123

Phone: (858) 966-1371

Fax: (858) 495-5841

*Contact: Maeve Hanley*

*Prepared by:*

**Conservation Biology Institute**

651 Cornish Drive

Encinitas, CA 92024

Phone: (760) 634-1590

Fax: (760) 634-1590

*Contact: Michael D. White, Ph.D.*

## TABLE OF CONTENTS

	<u>Page</u>
<b>ACKNOWLEDGEMENTS</b>	ii
<b>1.0 INTRODUCTION</b>	1
<b>2.0 STUDY AREA DESCRIPTION</b>	1
2.1 Onsite Land Uses	2
2.1.1 Ramona Airport	2
2.1.2 Ramona Municipal Water District	2
2.1.3 Cattle grazing	3
2.2 Geology and Soils	3
2.3 Climate	4
2.4 Hydrology	5
<b>3.0 METHODS</b>	5
3.1 Grasslands	5
3.1.1 Stephens' kangaroo rat	6
3.1.2 Grassland vegetation	8
3.1.3 Raptors	10
3.2 Vernal Pools, Vernal Swales, and Alkali Playas	11
3.2.1 Vernal pool, vernal swale, and alkali playa vegetation	11
3.2.2 Hydrology and water characteristics	12
3.2.3 Fauna	13
3.3 Santa Maria Creek Corridor	14
3.3.1 Water quality	14
3.3.2 Channel geomorphology	14
3.3.3 Riparian vegetation	14
3.3.4 Avifauna	15
3.3.5 Arroyo toad	16
3.4 Invasive, Nonnative Plant Removal	16
<b>4.0 RESULTS</b>	17
4.1 Grasslands	17
4.1.1 Stephens' kangaroo rat	17
4.1.2 Grassland vegetation	18
4.1.3 Raptors	20
4.2 Vernal Pools, Vernal Swales, and Alkali Playas	21
4.2.1 Vernal pool, vernal swale, and alkali playa vegetation	21
4.2.2 Hydrology and water characteristics	22
4.2.3 Fauna	23

	<u>Page</u>
4.3 Santa Maria Creek Corridor	24
4.3.1 Water quality	24
4.3.2 Channel geomorphology	25
4.3.3 Riparian vegetation	26
4.3.4 Avifauna	27
4.3.5 Arroyo toad	27
4.4 Invasive, Nonnative Plant Removal	28
5.0 DISCUSSION	28
6.0 REFERENCES	31

## **ATTACHMENTS**

ATTACHMENT A—PLANT SPECIES LIST

ATTACHMENT B—ANIMAL SPECIES LIST

ATTACHMENT C—PHOTOGRAPHS OF SELECTED SENSITIVE SPECIES

ATTACHMENT D—PHOTOMONITORING RESULTS

ATTACHMENT E—GRASSLAND SPECIES RICHNESS

ATTACHMENT F—VERNAL POOL SPECIES RICHNESS

ATTACHMENT G—WATER QUALITY RESULTS

## **LIST OF FIGURES**

Figure 1	Project location.	34
Figure 2	Conserved properties.	35
Figure 3	Soil series within the Preserve.	36
Figure 4a	Location of Preserve within the San Dieguito watershed.	37
Figure 4b	Estimated extent of the flooding of Santa Maria Creek in 2005.	38
Figure 5	Locations of field monitoring stations in the Preserve.	39
Figure 6	Density of SKR burrows and delineation of core SKR management areas in the Preserve.	40
Figure 7	Locations of rare plants and patches of purple needlegrass in the Preserve.	41

	<u>Page</u>
Figure 8	Percent cover of grassland species within categories of SKR habitat quality (i.e., low, medium, and high).
	42
Figure 9a	Average biomass ( $\pm$ SE) by SKR habitat quality.
	43
Figure 9b	Spring and fall 2006 biomass correlation.
	43
Figure 10	Average abundance of wintering raptors in the Preserve.
	44
Figure 11	Raptor species breeding in the Preserve.
	45
Figure 12	Vernal pool cover by pool zone in 2005 and 2006.
	46
Figure 13	Average depth of vernal pools by complex in 2005.
	47
Figure 14	Temperature, dissolved oxygen, and pH at monitoring stations SMC1, SMC2, and SMC3 for the 2005-2006 monitoring period.
	48
Figure 15	Stream flow and total suspended solids at monitoring stations SMC1, SMC2, and SMC3 for the 2005-2006 monitoring period.
	48
Figure 16	Total dissolved solids and conductivity at monitoring stations SMC1, SMC2, and SMC3 for the 2005-2006 monitoring period.
	49
Figure 17	Total nitrogen and total phosphorus at monitoring stations SMC1, SMC2, and SMC3 for the 2005-2006 monitoring period.
	49
Figure 18	Total coliform bacteria at monitoring stations SMC1, SMC2, and SMC3 for the 2005-2006 monitoring period.
	50
Figure 19	Chloride and sulfate at monitoring stations SMC1, SMC2, and SMC3 for the 2005-2006 monitoring period.
	50
Figure 20a	Channel geomorphology at cross-section SG2-SG6.
	51
Figure 20b	Channel geomorphology at cross-section SG7-SG11.
	52
Figure 21a	Tree and shrub cover at riparian transect monitoring locations in 2005.
	53
Figure 21b	Tree and shrub cover at riparian transect monitoring locations in 2006.
	54
Figure 22	Distribution of arroyo toads detected in the Preserve.
	55
Figure 23a.	Locations of invasive non-native plant species in the Preserve.
	56
Figure 23b	Locations of invasive non-native plant species in the Preserve.
	57
Figure 24	Aggressive non-native species treated in the Preserve.
	58

## LIST OF TABLES

	<u>Page</u>
Table 1      Grazing intensity (AUM) on Cagney Ranch, Oak Country Estates, and Davis-Eagle Ranch from 2004-2005.	4
Table 2      SKR distribution survey dates and areas surveyed.	6
Table 3      SKR burrow density classes as originally defined by O'Farrell (1992) and as scaled down for finer-resolution mapping in the field by S. Montgomery.	7
Table 4      Grassland plots and survey year.	9
Table 5      Grassland monitoring plots relative to SKR habitat quality rankings.	10
Table 6.      Geographic complexes supporting surveyed vernal pools, vernal swales, and alkali playas.	12
Table 7      Total number of seedlings recorded at each riparian transect location.	26

## APPENDICES (SEPARATE VOLUME)

- APPENDIX A—*Biological Survey Report for the Santa Maria Creek Restoration Project: Stephens' kangaroo rat* (Spencer and Montgomery 2006)
- APPENDIX B—*Wintering Raptors of the Cagney Ranch and Surrounding Ramona Grasslands (2003-2006)* (Wildlife Research Institute 2007)
- APPENDIX C—*Biological Survey Report for the Ramona Grasslands Preserve* (RECON 2005)
- APPENDIX D—*Biological Survey Report for the Santa Maria Creek Restoration Project: riparian birds* (Lovio 2006)
- APPENDIX E—*Biological Survey/Monitoring Report for the Santa Maria Creek Restoration Project: arroyo toads* (Hollingsworth et al. 2006)
- APPENDIX F—*Invasive Weed Report for the Santa Maria Creek Restoration Project: grassland and riparian invasive weed control efforts and results* (Kelly & Associates 2007)

## ACKNOWLEDGEMENTS

We would like to acknowledge the contributions of many organizations and individuals to the Ramona Grasslands Preserve Baseline Conditions Report. The following organizations and individuals conducted the technical studies that were the basis of the information reported herein: RECON Environmental, Inc., Technology Associates International Corporation (TAIC), Wildlife Research Institute (WRI), Kelly & Associates, John Lovio, San Diego Natural History Museum, Wayne Spencer (CBI), Steve Montgomery, Molly Pohl-Costello (SDSU), Fred Sproul, Gena Calcarone, and the City of San Diego Water Department. Susan Hector developed cultural resources information for the Preserve.

TAIC maintained the project database and prepared all of the maps used in this report. Gena Calcarone performed data entry and cataloged photographs. CBI staff Wayne Spencer, Nancy Staus, and John Bergquist contributed to data analysis for the project. Zachary Principe and Chris Basilevac with The Nature Conservancy provided invaluable field assistance and information on the properties owned by their organization.

We would like to thank Tellam & Tellam Cattle for their assistance with understanding their cattle operations in the Ramona Grasslands and their patience with the changes underway in the Preserve.

Finally, we would like to thank the California Water Resources Control Board for funding the Proposition 13 Grant for this project, Debbie Woodward with the San Diego Regional Water Resources Control Board for her assistance administering the Proposition 13 Grant, and staff at the San Diego County Department of Parks and Recreation, particularly Maeve Hanley, Jennifer Haines, and Carole Melum, for their help and contributions to the project.

## 1.0 INTRODUCTION

The Ramona Grasslands are located in the Santa Maria Valley, west of the town of Ramona in San Diego County (Figure 1). In 2003 the State Water Resources Control Board awarded a Proposition 13 grant to the County of San Diego for the protection and restoration of a portion of Santa Maria Creek and adjacent ephemeral aquatic habitats in the Ramona Grasslands Preserve (Preserve). The Preserve is comprised of a number of properties that have already been conserved or are publicly owned, properties with conservation agreements pending, and properties targeted for conservation (Figure 2). Additional land is anticipated to be included in the Preserve via landowner dedications as mitigation for future development projects (e.g., portions of Cumming Ranch). (Figure 2). The Preserve supports many of the unique biological resource values represented in the greater Preserve, provides a suite of important environmental services for the region, and embodies a rich cultural and historic heritage.

The Preserve hosts a unique assemblage of resources:

- The southernmost population of the federally endangered Stephens' kangaroo rat (*Dipodomys stephensi*);
- Unique vernal wetlands that support federally endangered San Diego fairy shrimp (*Branchinecta sandiegonensis*) and several rare plant species;
- An important population of the federally endangered arroyo toad (*Bufo californicus*) in Santa Maria Creek; and
- A diverse raptor community, including the largest population of wintering ferruginous hawks (*Buteo regalis*) in San Diego.

As part of the Proposition 13 grant project, a variety of field surveys and monitoring was implemented to characterize baseline conditions of the Preserve to inform future management and monitoring efforts. This Baseline Conditions Report summarizes the results of these surveys and monitoring efforts. Management and monitoring prescriptions are discussed in the Ramona Grasslands Preserve Area Specific Management Directives (CBI 2007).

## 2.0 STUDY AREA DESCRIPTION

The Ramona Grasslands are located in the Santa Maria Valley, situated between the coastal mesas and the mountains of the Peninsular Ranges in west-central San Diego County (Figure 1). The Santa Maria Valley is a broad basin (elevation 1,350-1,450 ft), surrounded by gentle hills and rocky rises vegetated with coastal sage scrub, chaparral, and oak woodlands. It lies within the Southern California Mountains and Valleys ecological section of the South Coast Ecoregion (Bailey et al. 1994, Goudey and Smith 1994, McNab and Avers 1994; Miles and Goudey 1998). The Preserve (Figure 2) is currently 1,950 acres in size; comprised of Davis-Eagle Ranch (1,230 acres), Cagney Ranch (420 acres), The Nature Conservancy (TNC) Oak Country Estates (230 acres), and Hardy Ranch (70 acres).

The core grasslands area of the Preserve remains relatively unfragmented, except by a few paved roads (e.g., Rangeland Road) and unpaved ranching roads (Figure 2). The Santa Maria Valley

itself is bordered by rural residential development and estate homes scattered on the hills surrounding the valley. South of the Preserve, houses line the south side of a portion of Santa Maria Creek along Voorhes Lane. Some areas around the periphery of the Preserve are used for dry farming and citrus and avocado orchards. Undeveloped land is located northwest, west, and east of the conserved lands. Northwest of the 1,950 acres of conserved lands is the 1,350-acre Gildred Family Ranch, which is currently under a purchase option with TNC. Adjacent to the conserved portion of Oak Country Estates (TNC Oak Country Estates) is the portion of Oak Country Estates, which has an approved residential development plan, and the Martz property, which is zoned as open space. East of the Cagney Ranch is Cumming Ranch, which currently has a proposed residential development plan, but a portion of which will likely be conserved as project mitigation or surplus open space land to be donated and/or sold to County of San Diego Department of Parks and Recreation (DPR). Northeast of Davis-Eagle Ranch is the proposed Montecito Ranch residential development (Figure 2). Offsite improvements, including the proposed Montecito Road, would accompany this development project. The proposed Montecito Road alignment extends south from Montecito Ranch, through the eastern portion of Davis-Eagle Ranch, and across Cumming Ranch.

## 2.1 Onsite Land Uses

### 2.1.1 Ramona Airport

The Ramona Airport lies at the eastern end of the Preserve, between Cagney Ranch and Davis-Eagle Ranch. The California Department of Forestry and Fire Protection has maintained the Ramona Air Attack Base at the airport since 1958. In 2002 the County extended the 4,000-foot runway an additional 1,000 feet to the west to accommodate larger fire-fighting aircraft. This extension, along with associated airport upgrades (e.g., sewer lines, taxiway, control tower), removed habitat occupied by the endangered Stephens' kangaroo rat (SKR), adversely affected vernal pools, and filled U.S. Army Corps of Engineers and California Department of Fish and Game jurisdictional wetlands and waters. As mitigation for these impacts, 62.5 acres of airport property were conserved (west and north of the extended runway in the western half of the airport property), 20.2 acres supporting vernal pools were conserved as part of the Preserve, and 1.34 acres of wetlands are to be created in the southwestern corner of Cagney Ranch within the Preserve. Habitat management plans for SKR (FAA 2002) and vernal pools (FAA 2003) were prepared to govern long-term management and monitoring of these target resources on the mitigation sites. Management of resources on the airport property will be performed by DPW, while long-term management of the resources in the Preserve (i.e., vernal pool and wetland mitigation sites, Figure 2) will be performed by DPR.

### 2.1.2 Ramona Municipal Water District

The Ramona Municipal Water District (RWMD) utilizes a portion of the Ramona Grasslands for storage and infiltration of treated sewage effluent. Treated effluent is piped from a treatment facility at the eastern end of the Ramona Grasslands to two storage reservoirs at the western end of the valley. Treated effluent is disposed via infiltration within a series of spray fields on the northwestern portion of Davis-Eagle Ranch. The irrigated spray fields are an important year-round source of green forage for cattle grazing of Cagney Ranch, Oak Country Estates, and

Davis-Eagle Ranch. Discussions are currently underway to determine if RMWD will acquire a portion of Davis-Eagle Ranch used for the spray fields or will enter into a long-term leases agreement with TNC for their use of the spray fields.

### 2.1.3 Cattle grazing

The majority of the core grasslands area has been used for cattle grazing for many years, with limited improvements such as perimeter fencing and wells installed over the years. Part of the Preserve (i.e., Cagney Ranch, TNC Oak Country Estates, and Davis-Eagle Ranch) is currently under lease to Tellam and Tellam Cattle for cattle ranching, while Hardy Ranch (and Cumming Ranch) is currently leased for grazing by Jack Dempsey. The Tellam and Tellam Cattle operation is the largest in the Preserve compared to Jack Dempsey, and consists of year-round cattle grazing, without formalized rotation or rest periods. Stocking rates are established on an annual basis, primarily based on weather and forage conditions (Tellam personal communication). No quantitative measures are made of forage production or residual dry matter. Bulls are added to the range around the first week of December to begin siring calves, with calving starting in mid-September of the following year. Calves are removed the following summer when the forage begins losing nutritional value. Supplemental feed is provided during summer (molasses supplement for increased protein and improved digestion of the dry forage), when the pregnant cows are on the range, which is otherwise low in nutrition once the vegetation dries out.

Grazing intensity, expressed as Animal Units per Month (AUMs) for the Cagney Ranch, Oak Country Estates, and Davis-Eagle Ranch is summarized in Table 2. DPR staff have estimated approximately 20 cows are currently grazing the Cumming and Hardy Ranch as a unit (McFedries personal communication). The lease agreement with Jack Dempsey allows a maximum of 70 cows to graze on Hardy Ranch.

## 2.2 Geology and Soils

The Santa Maria Valley basin is predominantly filled with soils of the Fallbrook and Bonsall series (Figure 3), which are well-drained to moderately well-drained sandy loams with a subsoil of clay loam or sandy clay loam over decomposed granodiorite, on gentle (2-9%) slopes (USDA 1973). On a more local scale, however, there is significant variation in soil characteristics depending on topographic location, depth of clay subsoils, and effects of erosion and deposition. Granodiorite outcrops dot the grasslands, predominantly on hilltops, with relatively deep, well-drained soils of decomposed granodiorites (in the Vista or Fallbrook series) sloping away from them. Lower-lying areas tend to support heavier clay soils, with shallow or even surface expression of clay hardpans. These soils sometimes develop characteristic vernal pool mima mound topography, which is best expressed on the Placentia soils in the Preserve near the southeast corner of the Ramona Airport. Gabbro outcrops and associated Las Posas soils are scattered through the grasslands likely influence plant associations (Sproul personal communication). Soils within the floodplain of Santa Maria Creek include deep, well-drained to excessively drained, sandy alluvium in the Visalia series (USDA 1973).

Soils have a strong influence on the distribution of target resources and management emphasis. Placentia soils have the greatest concentration of vernal pools. Bosanko clays dominate the low-lying eastern portion of the core grasslands and are present in patches on the western portion of the Preserve, and support existing native grasslands. Several sandy loams (e.g., Fallbrook and Vista) in the northern and western portion of the grasslands provide optimal habitat conditions for SKR. Soils along Santa Maria Creek are mostly sandy alluvial deposits—Tujunga sands along the stream channel and Visalia sandy loams in the adjacent floodplain. A series of alkali playas lies within areas mapped as Visalia sandy loams (USDA 1973), but these areas more likely have clay soil inclusions or eroded areas too small to have been mapped at the USDA mapping resolution.

**Table 1. Grazing intensity (AUM) on Cagney Ranch, Oak Country Estates and Davis-Eagle Ranch from 2004-2005.**

Month	2004 <sup>1</sup>	2005 <sup>2</sup>	2006 <sup>3</sup>
January	—	10/6	40/8/115
February	—	18/6	30/10/115
March	—	18/6	40/10/115
April	—	18/6	40/10/115
May	—	18/6	8/115 <sup>4</sup>
June	—	18/6	30/8/115
July	—	18/6	8/105 <sup>5</sup>
August	—	18/6	0/105 (135) <sup>6</sup>
September	20	18/6	—
October	20	18/6	—
November	20	18/6	—
December	20	18/6	—

— = No data

<sup>1</sup> AUM on Cagney Ranch

<sup>2</sup> AUM on Cagney Ranch/TNC Oak Country Estates

<sup>3</sup> AUM on Cagney Ranch/TNC Oak Country Estates/Davis-Eagle Ranch

<sup>4</sup> AUM on TNC Oak Country Estates/Cagney Ranch + Davis-Eagle Ranch, with 40 AUMs moved to the Vangler property this month.

<sup>5</sup> AUM on TNC Oak Country Estates/Cagney Ranch + Davis-Eagle Ranch

<sup>6</sup> AUM on TNC Oak Country Estates/Cagney Ranch + Davis-Eagle Ranch from 8/1 -8/28. Cagney Ranch + Davis-Eagle Ranch had 135 AUM from 8/28 – 8/31. No cows were on TNC Oak Country Estates during the month because of a mistakenly closed gate.

## 2.3 Climate

Temperature extremes at Ramona range from about 17°F to 112°F, with minimum mean temperatures in December-January of 37-38°F, and maximum mean temperatures during July-August of approximately 91°F (as recorded at the Ramona Airport). In the summer months the climate is generally hot and subhumid, with moderate oceanic influence. For the purposes of this report, rainfall data are summarized on a water year basis (October 1 – September 30), which is a common annual interval used by hydrologists in the U.S. to ensure annual stream flow and precipitation patterns are not split between calendar years. Rainfall is largely restricted to the period November through March, with 65% of the average annual rainfall of 15.86 inches/year falling from January to March. Inter-annual variation in rainfall can be high. For example,

during water years from 1974 to 2005, annual totals ranged from a minimum of 3.17 inches in 2001/2002 to a maximum of 34.08 inches in 1977/1978 (Western Regional Climate Center 2006). Rainfall totals for years 2002/2003, 2003/2004, and 2004/2005 were 17.38 inches, 7.64 inches, and 27.24 inches, respectively. Rainfall from October 1, 2005 through July 2006 is 7.94 inches (with reported March data incomplete).

## 2.4 Hydrology

The Ramona Grasslands comprise a significant portion of the Santa Maria Creek subbasin of the San Dieguito River watershed (Figure 4a). Santa Maria Creek and its tributaries drain about 57 mi<sup>2</sup> from the mountains east of Ramona, across the Ramona Grasslands, and through the steep and narrow walls of Bandy Canyon to its confluence with Santa Ysabel Creek. Below the confluence, the San Dieguito River flows through San Pasqual Valley into Lake Hodges, a City of San Diego drinking water reservoir listed as an impaired water body (Clean Water Act 303(d) listed) due to excessive nutrients and color from runoff of non-point source pollutants within the watershed.

Santa Maria Creek generally exhibits ephemeral flow in response to winter rainfall; although surface flow in the creek may persist very late in the year in heavy rainfall years and surface water is perennial at the far western end of the valley. The U.S. Geological Survey (USGS) has operated a stream gauge at the western end of the Santa Maria Valley (HUC 11028500), which measures all flow leaving the Ramona Grasslands. Annual mean daily streamflow of Santa Maria Creek from 1974-2005 was 3,968 cubic feet per second (cfs), and ranged from almost no flow for some years to 28,547 cfs in 1993. The Ramona Grasslands drain to Santa Maria Creek via ephemerally flowing swale features. During the extremely rainy 2004/2005, surface water left the Santa Maria Creek channel near the western boundary of Hardy Ranch (peak flow 2,050 cubic feet per second). Surface water flowed through an east-west trending swale across the grasslands, where it crossed Rangeland Road and then re-entered Santa Maria Creek via a tributary swale (Figure 4b). This east-west trending swale supports the alkali playa system.

## 3.0 METHODS

Resource-specific quantitative surveys were conducted in 2005 and 2006 for the Preserve and are described in the sections below. In addition, qualitative botanical surveys were conducted in 2005 and 2006. These surveys focused on mapping occurrences of sensitive plant species encountered in the Preserve as other field activities were conducted. The results of these qualitative surveys are presented in Section 4.

### 3.1 Grasslands

Surveys were conducted in the grassland habitats of the Preserve to map habitat quality for Stephens' kangaroo rat, assess vegetation composition and structure, and use of the preserve by raptors for wintering and breeding.

### 3.1.1 Stephens' kangaroo rat

Detailed SKR survey methodologies can be found in *Biological Survey Report for the Santa Maria Creek Restoration Project: Stephens' kangaroo rat* (Spencer and Montgomery 2007, Appendix A), and are summarized below.

A primary aim of this study was to create a comprehensive SKR distribution map for the Preserve to inform future management and monitoring actions. This was accomplished by surveying all properties within the Preserve and supplementing these surveys with existing information from properties adjacent to the Preserve, most notably the Ramona Airport. Some adjacent properties are also known from previous surveys not to support SKR or suitable habitat (e.g., Cumming Ranch; O'Farrell 2000a, 2004). SKR absence was inferred for some properties for which access was not granted (e.g., Hobbs and the "Voorhes Lane properties") based on lack of suitable habitat, as indicated by inspection of aerial photographs and ground-truthing from property boundaries. However, no attempt was made to map SKR distribution on some properties that are known to support SKR based on previous trapping surveys (P. Vergne, unpublished data), but that could not be confidently map without access (e.g., Martz and RWMD ownerships).

On properties within the Preserve, distribution and relative abundance of SKR were mapped in the field by Wayne Spencer and Stephen Montgomery, with assistance from Esther Rubin and Scott Tremor (Table 1). During 2005, surveys covered properties included in the Preserve at that time (Cagney, Hardy, and TNC Oak Country Estates). Davis-Eagle Ranch was added to the reserve area in December 2005 and surveyed for SKR during 2006. During 2006 a number of areas previously surveyed during 2005 were spot-checked to confirm that SKR distribution had not changed notably from one year to the next, so the composite 2005-2006 map could be treated as one consistent baseline data source.

**Table 2. SKR distribution survey dates and areas surveyed.**

Date	Observers*	Area Surveyed
2005		
23-Sep	WS, SJM	North and west TNC Oak Country Estates
12-Nov	WS, SJM	East end Cagney, Hardy
19-Nov	WS, SJM	Central Cagney
17-Dec	WS, ST	South Cagney, south and central TNC Oak Country Estates
2006		
25-Aug	WS	Southwest Eagle Ranch and spot checks on Cagney
5-Sep	WS, SJM, ER	Central and north Davis-Eagle Ranch and spot checks on TNC Oak Country Estates
6-Sep	WS, SJM, ER	North and northeast Davis-Eagle Ranch
8-Sep	WS, SJM, ER	South and central Davis-Eagle Ranch and spot checks
26-Oct	WS	West-central Davis-Eagle Ranch and spot checks on airport, Cagney
26-Oct	WS	Northwest Davis-Eagle Ranch and spot checks on Cagney

\* WS = Wayne Spencer, SJM = Stephen Montgomery, ST = Scott Tremor and ER = Esther Rubin

The mapping method involved walking meandering transects over the entire area (at no greater than 50-m spacing) searching for signs of SKR occupancy (burrows, scats, tracks, dust baths). Once signs of occupancy were found in a particular location, the biologists searched for the outer perimeter of the occupied area (where no further sign could be found, or where habitat clearly became unsuitable), enclosed it with a polygon, and classified the relative density of SKR burrows within the polygon using density classes originally developed by Michael O'Farrell (1992) and modified by Stephen Montgomery for ease in mapping at finer resolution (Table 3). Results were marked onto 1:3200-scale, true-color aerial photographs. Mapping was aided by having the aerials gridded with 50 x 50-m cells and by use of GPS. During 2005 and 2006 surveys of the Preserve, most occupied habitat supported only trace SKR densities; moderate-density areas were very rare, and there were no high density areas to map.

**Table 3. SKR burrow density classes as originally defined by O'Farrell (1992) and as scaled down for finer-resolution mapping in the field by S. Montgomery.**

Density Class	Burrows/ha (O'Farrell)	Burrows/200 m <sup>2</sup> (Montgomery)
Trace	<50	<1
Low	50-200	1-4
Moderate	200-700	4-14
High	>700	>14

SKR density was also mapped at the edges of the Ramona Airport, and incorporated an edge-matched distribution and density mapping performed on the Airport property in 2005 by Haas and O'Farrell (2005). The Haas and O'Farrell (2005) polygons were converted to a similar mapping resolution as the current data and applied the same density classes as were done on Preserve. The polygons near the Airport boundary were calibrated and adjusted as necessary based on the current observations. In addition to survey dates listed in Table 2, which all reflect ideal sign-survey conditions during late summer-fall, Wayne Spencer also spot-checked portions of the study area during winter-spring conditions on January 26 and April 28, 2006.

Limited trapping surveys were performed in portions of the Preserve to confirm which species of kangaroo rat was present, SKR or the non-listed Dulzura kangaroo rat (*Dipodomys simulans*; DKR; formerly *D. agilis*). Although these two species sometimes co-occur at a local scale, SKR are competitively dominant and almost always occupy the most open grassland habitats, whereas DKR are generally restricted to those areas with some scrub cover (Price et al. 1991). Previous intensive trapping surveys in the Ramona Grasslands (e.g., Ogden 1998, Spencer 2002, P. Vergne unpublished data) have repeatedly reinforced these observations, with only SKR found in the open grasslands but either species occurring grass/scrub interface areas and predominantly DKR in open scrub habitats or oak savannahs. Consequently, sample-trapping was conducted to identify which of the two species was present in scrub interface areas and refine the mapping of SKR-occupied habitat areas.

### 3.1.2 Grassland vegetation

#### *Field surveys*

Quantitative vegetation surveys of the Preserve were conducted in 2005 and 2006, and included determination of species composition, vegetative cover (including amount of bare ground and thatch), plant height, and biomass. Plot locations varied over the two year period as additional properties were added to the Preserve. Plot numbers and survey year are summarized in Table 4 and shown in Figure 5. All field surveys were conducted by DPR staff Fred Sproul and Gena Calcarone.

Grassland surveys were conducted from May to July each year using a point-intercept methodology. In 2005, two 50-meter transects were established perpendicular to each other at each plot location. In 2006, this was changed to a single 100-meter transect at each plot location. At every meter along each transect a pin was dropped, and every species (or bare ground or thatch) touching the pin and their height was recorded. In 2006, disturbance factors were recorded at each pin location, if present, and the grazing intensity across each plot (low, medium, high) was noted. In addition, using the same 100 meter transects, a two meter wide belt transect was established at each plot location. Every plant species within the belt transect was recorded.

Biomass samples were collected in October 2005, January 2006, May 2006, and October 2006. At each plot location, between two and four (typically three) biomass samples were collected. Biomass samples were obtained by collecting all above-ground biomass, including rooted vegetation and thatch, from within a 13.25-inch interior diameter hoop (hoop area is 0.96 feet<sup>2</sup>). The collected vegetative material was air-dried and weighed. The weight in grams was converted to pounds per acre (lbs/acre) by multiplying by 100. For the purposes of this document, biomass refers to vegetative material collected in the spring at peak growth (May samples) and Residual Dry Matter (RDM) is material collected in the fall and early winter when annual plants are dead (October and January samples). Thus, RDM represents the vegetative material remaining in the grasslands prior to the following year's new growth.

The native bunchgrass, purple needlegrass (*Nasella pulchra*), was present in portions of the Preserve. At each plot, all individual purple needlegrass bunches within a 2-m wide x 100 m belt transect were counted.

Photo-monitoring was conducted in both 2005 and 2006. Photographs were taken at monitoring plot 14 in each of the four cardinal directions.

**Table 4. Grassland plots and survey year.**

<b>Plot Number</b>	<b>2005 Vegetation and Biomass</b>	<b>2006 Vegetation and Biomass</b>	<b>2005 Biomass Only</b>
1		X	
2	X		
3	X	X	
4	X	X	
5	X	X	
6	X	X	
7	X	X	
8	X		
9	X	X	
10	X	X	
11	X	X	
12		X	
13		X	
14	X	X	
15		X	
16	X	X	
17	X	X	
18	X	X	
19		X	
20	X	X	
21		X	
22		X	
23		X	
24		X	
30			X
31			X
32			X
33			X
34			X
35			X
36			X
37			ND
38			X

ND = No data collected

### *Data analysis*

For the purposes of this report, survey results were generally summarized by geographic areas supporting differential habitat quality for Stephens' kangaroo rat, as discussed in Section 3.1.1. Table 5 shows the distribution of grassland monitoring plots among SKR habitat quality

categories. For all vegetation metrics (e.g., percent cover, average height, or biomass), a single value was derived for each plot and then a mean value calculated for each Stephens' kangaroo rat habitat quality category by averaging the plots within that category. For data collected with the point intercept methodology, absolute percent cover for each species (or bare ground or thatch) was calculated as the number of points that hit that species divided by the total number of points. Average height was calculated as the average of all height measurements recorded for each species at each plot. Similarly, average biomass within each plot was the average of the replicate samples collected at each plot.

**Table 5. Grassland monitoring plots relative to SKR habitat quality rankings.**

**Note: there are no grassland monitoring plots 24-29.**

Plot Number	Habitat Quality	Plot Number	Habitat Quality	Plot Number	Habitat Quality
1	Low	12	Low	23	Low
2	Low	13	Low	24	Low
3	Medium	14	Low	30	High
4	Low	15	High	31	Low
5	Low	16	Low	32	High
6	Low	17	Low	33	High
7	Medium	18	Low	34	High
8	Medium	19	Low	35	Medium
9	Medium	20	Low	36	High
10	High	21	Medium	37	Medium
11	High	22	High	38	Low

### 3.1.3 Raptors

Detailed raptor survey methodologies can be found in *Wintering Raptors of the Cagney Ranch and Surrounding Ramona Grasslands (2003-2006)* (Wildlife Research Institute [WRI] 2007, Appendix B), and are summarized below.

#### *Historical Data*

Raptor observational data were compiled from WRI's winter Hawk Watch in the Preserve, held January through February in 2003, 2004, 2005, and 2006. These observations focus primarily on the grasslands around the WRI property and along Rangeland Road, just north of WRI, with supplemental observations north of Voorhes Lane and surrounding the Ramona Airport (Figure 2). Wintering raptor survey sites are shown in Figure 5. In 2003, 2004, and 2005 only sites 1-3 were surveyed, and all three of these years contain some weeks in which two days of observation data were collected. Few documented observations were made at sites 4-7 during these years. For 2003 only, observations started in December and covered the time frame between December 28, 2002 and February 2, 2003. Although observations were made for ten weeks in 2005, written documentation for that year is limited.

### *Current Data Collected (Year 2006)*

Data specific to the Baseline Conditions project was collected during January and February 2006, when migrating raptors are most likely to be in the area and all raptors are more visible due to decreased foliage in trees along Santa Maria Creek. Surveys were performed one day per week from January 1 through February 28, 2006 at monitoring sites 1-3 (Figure 5). Between January 14 and January 28, 2006, data were collected for three days at all sites.

Observations were made from 0900 to 1200 hours at sites 1-3, with approximately one hour spent observing at each site. The data from sites 4-7 were collected on three separate surveys conducted between January 14 and January 28, 2006. The surveys were conducted for two hours between 1200 and 1400. A total of fifteen minutes of observations were collected at each of the three sites. Multiple observers performed the initial spotting of raptors and one experienced raptor biologist identified and recorded observations for individual species. Observations were recorded on a standard observation form. Observations were made with Kowa 10 x 42 binoculars and 10 x 20-60 zoom scopes were used in raptor identification.

### *Data Analysis*

The numbers of wintering raptor surveys varied greatly among years. Numbers of surveys by year were: 2003 = 10, 2004 = 8, 2005 = 4, and 2006 = 13 (WRI 2007, Appendix B). Therefore, for purposes of this report, the mean number of raptors for each survey year was calculated by averaging the count of each raptor species for all surveys during a given year. All nesting locations were mapped in the field.

## 3.2 Vernal Pools, Vernal Swales, and Alkali Playas

Monitoring was conducted on ephemeral aquatic habitats in the Preserve to characterize hydrology and water characteristics, faunal communities, and composition of vegetation communities. Ephemeral aquatic habitats in the preserve include vernal pools, vernal swales, and alkali playas. Vernal swales are natural linear depressions that are part of the surface drainage network of Santa Maria Creek, and periodically pond and function like vernal pools. Vernal pools, vernal swales, and alkali playas monitored in the Preserve are shown in Figure 5. Monitoring in 2005 did not start until January, after pools began filling in October of 2004. Thus, some of the information on pool hydrology and fauna may not accurately represent the true conditions at the monitored locations.

### 3.2.1 Vernal pool, vernal swale, and alkali playa vegetation

#### *Field surveys*

Quantitative vegetation surveys of the vernal pools were conducted from April – July 2005 and during May 2006. At each pool, one transect was established on randomly selected compass bearing and a second established perpendicular to the first. Ten, 10 cm x 50 cm quadrats were randomly selected across these transects, generally five quadrats to each transect. Each quadrat was assigned to a visually estimated depth zone within each pool: 1) center (deepest area),

2) intermediate, and 3) pool edge (upland transition). The cover of all plant species (including bare ground and thatch) was recorded within each quadrat. In 2006, any disturbances associated with each quadrat were recorded, and grazing intensity (low, medium, high) was visually estimated. All species found in a pool, whether or not present in a quadrat, was separately recorded for each vernal pool. All field surveys were conducted by DPR staff Fred Sproul and Gena Calcarone.

Photo-monitoring was conducted in both 2005 and 2006. Photographs were taken at vernal pools e44, e45, e53, and e54 in each of the four cardinal directions.

### *Data analysis*

For purposes of this report, vegetation data are summarized as percent absolute cover by averaging the quadrats within each depth zone for each pool. Thus, species-specific percent cover estimates are calculated separately for each zone in every pool, and pools serve as the sampling unit for summarizing vegetation data. Pools, swales, and alkali playas were grouped into geographic complexes for purposes of data summary and presentation. These surveyed pools, and the complexes to which they were assigned, are summarized in Table 6. The locations of individual surveyed pools are shown in Figure 5.

**Table 6. Geographic complexes supporting surveyed vernal pools, vernal swales, and alkali playas.**

<b>Airport Vernal Pools</b>	<b>Cumming Vernal Swale</b>	<b>Cagney Swale</b>	<b>Cagney Vernal Pools</b>	<b>Alkali Playa</b>
e44	ev1	vs1	e56	RAAP100
e45	ev2	vs2	e58	
e46		vs3	e59	
e52		vs4	e62	
e53				
e54				
e77				
r24				

### 3.2.2 Hydrology and water characteristics

Hydrology and water characteristics methodologies for vernal pools can be found in *Biological Survey Report for the Ramona Grasslands Preserve* (RECON 2005, Appendix C), and are summarized below.

In January 2005, a staff gauge was installed in each of the study vernal pools. Staff gauges were constructed of 24-inch-long sections of 1-inch diameter polyvinyl chloride (PVC) pipe, fitted over 30-inch-long sections of 1/2-inch diameter rebar. The gauge was fitted so the PVC pipe and rebar were flush on one end, and the flush end was capped to indicate the top of the gauge. From the base of each cap, marks were placed at 1-inch increments to indicate depth. Self-adhesive numbers indicating the pool's identity were placed vertically down the gauge, opposite the side with the 1-inch hash marks. The staff gauge was then driven into the ground so that the bottom

hash mark, indicating zero inches, was flush with the ground in the deepest portion of each pool. Water depth was monitored weekly from January 21, 2005 to April 19, 2005. Water depth to an accuracy of 0.25 inch was recorded for each vernal pool on each date. For purposes of analysis and presentation in this report, water depth was averaged across vernal pools within complexes shown in Table 6.

Dissolved oxygen levels in each vernal pool were monitored monthly using a LaMotte Dissolved Oxygen Test Kit (Code 7414/5860). Dissolved oxygen testing followed instructions and procedures as outlined in the Dissolved Oxygen Test Kit manual. Water temperatures were recorded every two weeks. In addition, anecdotal information on conductivity (a measure of the amount of dissolved solids, such as salts, in water) was collected in several alkali playas and several vernal pools in the Airport complex in March 2005.

### 3.2.3 Fauna

Methodologies for vernal pool faunal surveys can be found in *Biological Survey Report for the Ramona Grasslands Preserve* (RECON 2005, Appendix C), and are summarized below.

#### *Fairy shrimp*

Fairy shrimp surveys were conducted according to U.S. Fish and Wildlife Service (USFWS) survey guidelines (USFWS 1996) every two weeks, starting January 21, 2005 and ending April 1, 2005. During each survey, the following steps were followed at each pool:

1. Prior to disrupting the water surface of the pool, if the view was relatively clear and unobstructed, the surveyor examined the pool for fairy shrimp to estimate the number of shrimp present.
2. Air temperature, water temperature, and maximum water depth [using staff gauges (see Section 3.2.1 above)] were recorded.
3. Using an aquarium fish net attached to an extendable painters pole, the surveyor made three-foot-long sweeps through the water to catch any fairy shrimp or other aquatic species that may have been present. All species caught in the net were examined, identified, and then returned to the pool; except fairy shrimp samples that were collected for identification and accessioning.
4. Step 3 was repeated in different locations around the vernal pool approximately 15 to 30 times depending on the size of the vernal pool.

#### *Vernal pool amphibians*

Amphibian surveys focused on species that occur in vernal pools such as western spadefoot toad (*Spea hammondi*), Pacific tree frog (*Pseudacris regilla*), and western toad (*Bufo boreas*). These surveys were conducted every two weeks, between January 21 and April 1, concurrent within fairy shrimp surveys. At each pool, the presence or absence, estimated number (e.g. 10s, 100s,

1000s, etc.), and the lifecycle stage (e.g. egg cluster, tadpole, or toadlet) of each species was noted.

### 3.3 Santa Maria Creek Corridor

Surveys were conducted in the Santa Maria Creek corridor to characterize water quality, channel geomorphology, vegetation composition and structure, breeding season avifauna, and the distribution of arroyo toads.

#### 3.3.1 Water quality

Water quality monitoring in Santa Maria Creek was conducted by the City of San Diego Water Department. The water quality monitoring program established three sampling locations: SMC1 is located immediately upstream of the Ramona Grasslands, SMC3 is located in the central portion of Santa Maria Creek in the project area and is within a reach of the creek traversing privately owned residential properties, and SMC2 is located immediately east of the Rangeland Road bridge (Figure 5). Santa Maria Creek was sampled twice monthly during 2005 and early 2006, when stream flow was present. Sampling was conducted per the Santa Maria Creek Restoration Water Quality Monitoring Plan, prepared by the City of San Diego Water Quality Laboratory (2004). Temperature, pH, oxidation-reduction potential (ORP), dissolved oxygen (DO) and specific conductance, and stream flow were measured in the field. All other chemical parameters were analyzed in the City of San Diego Water Quality Laboratory.

#### 3.3.2 Channel geomorphology

The channel geomorphology of Santa Maria Creek was measured at ten cross sections in the Preserve (Figure 5). Vertical elevation-horizontal distance pairs were measured in the field using survey equipment, providing the cross-sectional shape of the channel, while GPS coordinates provided the latitude and longitude of the benchmarks (right and left end points of the cross-sections). Approximate bankfull geometry for each cross-section in the Santa Maria Creek reach was established from standard field observations and measurements. All cross-sections were standardized to read from river left (RL) to river right (RR) when the observer is looking downstream. Channel geomorphology is presented in real elevations (meters above mean sea level). All field surveys were conducted by Molly Pohl-Costello and Michael White.

#### 3.3.3 Riparian vegetation

At each cross-section location (Figure 5), riparian vegetation community composition and structure was measured using a line-intercept methodology. At each location, a transect line was established across the creek channel. The intercept lengths of individual plant species in the canopy layer, shrub layer, and herb layer along each transect was recorded. In 2006, the herb layer was quantified using a point-intercept methodology (i.e., recording the species touching a pin dropped every one-half meter along the prescribed transect). In addition, all riparian shrub and tree species were counted within a 2-meter belt transect at each location and categorized as seedlings or adults. All field surveys were conducted by DPR staff Fred Sproul and Gena Calcarone.

Photo-monitoring was conducted in both 2005 and 2006. Photographs were taken at each location.

#### 3.3.4 Avifauna

Detailed survey methodologies for riparian avifauna can be found in *Biological Survey Report for the Santa Maria Creek Restoration Project: riparian birds* (Lovio 2007, Appendix D), and are summarized below.

A breeding-bird census (Van Velzen 1972) was conducted for riparian vegetation along Santa Maria Creek, utilizing a technique known as “spot-mapping” (Bibby et al. 1992, Ralph et al. 1993). Under the assumption that pairs of breeding birds occupy regular areas that are at least partially exclusive of other pairs of the same species during the breeding season and that territorial birds advertise their presence by visual and auditory clues, census areas are completely and systematically surveyed on multiple visits during a single breeding season. The breeding-bird census was conducted on seven dates over a period of 37 days between mid-May and mid-June, 2005. During each census visit, the locations and behaviors of individuals of all species detected were recorded on a map of the census area. Over repeated visits, the cumulative map registrations for each species tended to form distinct clusters that represent different pairs of a given species.

Daily census visits were begun shortly after dawn to maximize the use of higher morning bird activity. Starting points on the creek and directions of movement were varied among the census visits, such that each section of the creek was covered at various times of day throughout the census period in an effort to minimize bias from differences in bird activity attributable to time of day. Bird locations for all species were marked on a separate high-resolution aerial photograph for each visit and associated demographic and behavioral data were recorded on a standard data form.

Interpretation of summary maps for the various species involved two somewhat overlapping steps: 1) Initial recognition of map clusters that likely represent separate breeding pairs or other units of the species. This step employed several basic criteria for qualifying any group of map registrations as a potential breeding unit of a species: a) Some level of obvious clustering of registrations relative to the overall dispersion of registrations for the species throughout the study area; criteria for clustering accounted for the scale of movement (generally the reciprocal of density) of the particular species; b) inclusion of registrations from a minimum of three dates spanning at least two weeks (approximately half of the 37-day span of the census period); c) presence of a nest or other definitive evidence of nesting if criteria a or b were lacking or insufficient. 2) Separation of clusters from adjacent clusters of the same species. Clues involved in this process included: a) gaps between clusters in otherwise continuous habitat; b) simultaneous or nearly simultaneous territorial displays by adjacent pairs; c) counterpart territorial registrations close in time in each cluster on one or more dates (greater confidence of distinctness of clusters with more dates).

The bird census was conducted uniformly along the entire study area, irrespective of habitat types and political boundaries. However, in the locational data analysis, map registrations and breeding territory cores (clusters) were recorded as occurring within any of six stream reaches (A-F) that correspond to property boundaries (Figure 5). Habitat within each of these segments is fairly uniform as a result of natural and anthropogenic factors and, with the exception of Reach A, the lengths of the segments are roughly comparable. The delineation of these segments, and the categorization of bird data within them, provides for simultaneous avifaunal comparisons among the habitat types.

### 3.3.5 Arroyo toad

Detailed methodologies for arroyo toad surveys conducted in 2005 can be found in *Biological Survey Report for the Ramona Grasslands Preserve* (RECON 2005, Appendix C). Detailed methodologies for arroyo toad surveys conducted in 2006 can be found in *Biological Survey/Monitoring Report for the Santa Maria Creek Restoration Project: arroyo toads* (Hollingsworth et al. 2006, Appendix E). These methods are summarized below.

Day and nighttime directed sight surveys for arroyo toads were conducted along Santa Maria Creek with the TNC Oak Country Estates portion of the Preserve between March 16 and June 14, 2005, and the entire Preserve on May 31 and June 12, 2006. These surveys followed the guidelines of the U.S. Fish and Wildlife Service (1999) and in 2006 were augmented with recommendations from United States Geological Survey (USGS 2005). On each survey date, two to three biologists walked along the edge or within the creek to detect the presence or absence of arroyo toads. Surveys were confined to the main channel of the creek bed; upland habitats beyond the banks of stream channel were not surveyed.

Data were recorded on datasheets and in field notebooks to document life stage, time, location, habitat, air and water/substrate temperatures, and signs of disturbance. In 2006, photographic vouchers were recorded for all sight records provided the animal's position allowed for photography, and locations were recorded with a handheld Garmin Legend GPS unit using an accuracy reading of six meters or less. A handheld Coleman lantern, Canon high intensity video light, and headlamps assisted nighttime surveys.

### 3.4 Invasive Non-native Plant Removal

The locations of invasive non-native plant species in the Preserve were initially identified by TNC in 2004, and included artichoke thistle, intermediate wheatgrass, giant reed, and salt cedar. New locations of these species were identified during removal efforts in 2005 and 2006 and were treated when detected. Previously unmapped invasive non-native plants species were detected in the Preserve during 2005 and 2006 removal efforts, and included milk thistle, Italian thistle, horehound, and perennial pepperweed (RECON 2005 and Kelly & Associates 2007, Appendices C and F). Invasive non-native plants were found throughout the Preserve properties, often times adjacent to sensitive or rare biological resources. Due to the sensitive nature of many flora and fauna in the Preserve, extra care was taken when selecting the methodology for removal of invasive non-native plants. Invasive non-native plants were treated repeatedly through 2005 and 2006, and an additional treatment is anticipated during the winter of 2007.

A mixture of hand removal, and herbicide use was employed to accomplish the invasive non-native plant removal objectives. Herbicide was employed most frequently, and a 50-gallon truck sprayer was used when dense concentrations of plants occurred, while backpack sprayers were used in less dense situations. An alternative herbicide application method was the “cut stump” technique used on arundo and saltcedar. In the cut stump technique, the saltcedar or arundo plant is cut with loppers, chainsaw, or handsaw, then a concentrated herbicide is applied to the cut stump within one minute of cutting. Herbicides used for invasive plant removal included Transline® and AquaMaster™. Transline®, clopyralid, Garlon 4a, Pathfinder (a pre-mix of Garlon 4a in a seed oil base), Glypro Pro (a Glyphosate herbicide, a generic Roundup), and Fusilade II. See *Biological Survey Report for the Ramona Grasslands Preserve* (RECON 2005, Appendix C) and *Invasive Weed Report for the Santa Maria Creek Restoration Project: grassland and riparian invasive weed control efforts and results* (Kelly & Associates 2007, Appendix F) for details of herbicide applications.

## 4.0 RESULTS

Lists of plant and animal species detected in the Preserve, including their sensitivity status, are provided in Attachments A and B respectively. Photographs of selected sensitive species are provided in Attachment C

### 4.1 Grasslands

#### 4.1.1 Stephens' kangaroo rat

Detailed results SKR surveys are provided in *Biological Survey Report for the Santa Maria Creek Restoration Project: Stephens' kangaroo rat*, (Spencer and Montgomery 2007, Appendix A), and are summarized below. SKR is federally endangered, state threatened, and a County of San Diego Group 1 species.

The majority of suitable and occupied SKR habitat is distributed in a broad, arcing mosaic of mostly well-drained, hilly topography near the center of the grasslands, with smaller mosaics or isolated pockets of suitable habitat scattered in other areas (Figure 6). The largest, most contiguous concentration curves around the west end of the Ramona Airport and extends west to Rangeland Road in those areas not used as effluent spray fields.

A second concentration of SKR habitat occurs in association with the northern fringe of the grasslands, where hills supporting coastal sage scrub rise up from the grasslands on the northern portion of Davis-Eagle Ranch. Both SKR and DKR were captured in this northern fringe area (Figure 6), with SKR occurring in the more open or down-slope portions, and DKR more in the edges of the coastal sage scrub and along a dirt road through sage scrub. Some habitat polygons found to have sign of kangaroo rats in this area were therefore omitted from SKR density polygons in Figure 6, as it was concluded they were unlikely to support SKR and highly likely to support DKR.

Smaller and more isolated pockets of habitat are found outside these two primary concentrations or core areas of habitat. On TNC Oak Country Estates, SKR was captured on a broad sandy flood plain near Santa Maria Creek, which was mapped as occupied at trace densities (although it may qualify as occupied at low densities). Several small pockets of trace or potential SKR habitat was mapped on and around isolated rocky hills on TNC Oak Country Estates, SKR presence was confirmed during previous trapping surveys (O'Farrell 2000b, 2002). Other isolated pockets of trace-occupied or potential habitat are also associated with rocky hills rising out of less suitable clay soils on portions of Cagney and Davis-Eagle Ranch, including some between the RMWD effluent spray fields.

Most areas mapped as unsuitable for SKR consist of heavier clay soils, such as eastern portions of Cagney and Davis-Eagle Ranch, much of TNC Oak Country Estates, and all of the Hardy and Cumming properties. Heavier clay soils also separate the large mosaic of habitat in the middle of the grasslands from the occupied areas along the northern fringe. Loose alluvial soils in the floodplain of Santa Maria Creek in the southern part of Cagney Ranch are also not occupied by SKR. This may be attributed to one or more of the following hypotheses: (1) these very loose, sandy soils may not be able to sustain SKR burrows, which may collapse easily in them; (2) occasional flooding by Santa Maria Creek may eliminate SKR from the area (drowning, wetting, and displacement); and (3) denser than average growth of annual grasses and associated thatch, perhaps due to lesser grazing intensity or elevated ground water.

The creation of the effluent spray fields in the western portions of Davis-Eagle Ranch may have apparently rendered some previously suitable habitat unsuitable, due to saturation of the soil and creation of dense, irrigated vegetation. Previously the mosaic of occupied habitat patches was probably more contiguous through this area. Although a few pockets of well-drained soils between the sprayfields are currently occupied, and more areas are probably occupied in years of expanded SKR populations, for the most part the interstices between spray fields appear to be somewhat degraded in habitat quality due to drifting spray, which elevates soil moisture relative to natural conditions. Only the larger and better drained rises between spray fields are likely to reliably support SKR from year to year.

Considered together, all these observations indicated that SKR populations were very low but relatively stable over the survey period, with little evidence of population expansion or contraction during 2005-2006 (but following a dramatic contraction from 2004 to 2005; Haas and O'Farrell 2005). The winter of 2004-2005 was the wettest on record at the Ramona Airport (29.03 inches of rain), which led to extraordinary growth of grasses during 2005. The SKR population contracted in response to this change in vegetation, with SKR persisting only in the most well-drained and highly suitable soils. The population did not appear to expand significantly in the drier conditions of 2006.

#### 4.1.2 Grassland vegetation

A variety of rare plant species were detected in the Preserve (Figure 7), including:

San Diego thornmint (*Acanthomintha ilicifolia*); FT, SE, County Group A  
Little barley (*Hordeum intercedens*); County Group C

Small-leaved morning-glory (*Convolvulus simulans*); County Group D  
Round-leaved filaree (*California macrophylla*); County Group B  
Southern tarplant (*Centromadia parryi* var. *australis*); County Group A  
Graceful tarplant (*Holocarpha virgata* ssp. *elongata*); County Group D  
(FT = federally threatened, SE = California Endangered)

The majority of these species are associated with clay soils, which are generally found around the margins of the Preserve (Figure 7). Small-leaved morning-glory, round-leaved filaree, and San Diego thornmint are only found in the eastern portion of the Preserve on the flanks of the knoll on the Hardy Ranch property. Purple needlegrass is scattered on clay soils throughout the Preserve, where it is often associated with graceful tarplant, but is particularly abundant on the southern portion of TNC Oak Country Estates. Little barley is known at one location in the eastern-central portion of the Preserve. Southern tarplant is abundant in the east-west trending swale north of Santa Maria Creek.

### *Species richness*

The grasslands support a moderately rich assemblage of plant species, although a small number of these species were most frequently detected in grassland plots and these are largely non-natives. Grassland plots with the highest number of species (> 30 species per plot) were generally associated with rocky knolls (e.g., Plot #3, 9, 14, and 19; Figure 5 and Attachment E). A total of 110 taxa were recorded in the grassland plots in 2005 and 2006 (Attachment E). However, 12 of these taxa (highlighted in Attachment E) were found in 16 or more of the 24 grassland plots, representing an average frequency of 5% of the total species in each grassland plot. Of these 12 frequently occurring species, only 3 of these taxa are native to the grasslands. Sixty seven of the 110 taxa were only found in 5 or fewer plots, representing an average frequency of 1.4% of the total number of species in each plot.

### *Structure and composition of grasslands*

Cover in the grasslands tended to be dominated by non-native annual grasses and forbs, primarily slender wild oat (*Avena barbata*), ripgut grass (*Bromus diandrus*), long-beaked filaree (*Erodium botrys*), Italian ryegrass (*Lolium multiflorum*), common catchfly (*Silene gallica*), and hairy rat-tail fescue (*Vulpia myuros* var. *hirsuta*) (Figure 8). The structure and composition of grassland habitats tended to vary among the three categories of SKR habitat quality. While no statistical differences were detected, the amount of bare ground, the cover of forbs, particularly from *Erodium* spp., and the forb to grass ratio tended to be higher in high quality habitat. Higher cover of annual grasses and thatch were observed in lower quality habitat.

The exceedingly wet fall and winter of 2004-2005 produced dense growth of grassland vegetation. Vegetative cover was higher across all SKR habitat quality categories in 2005, with the cover of slender wild oat, long-beaked filaree, and Italian ryegrass particularly high (Figure 8). In 2006, these species, along with ripgut grass, provided the highest contributions to grassland cover, but at lower levels than observed in 2005.

Although statistical differences were not found, RDM in 2006 was on average lower in high quality SKR habitat than in either medium or low quality SKR habitat (Figure 9a). High quality SKR habitat had a mean RDM of approximately 1,800 lbs/acre, whereas medium and low quality habitat had mean RDMs over 2,300 lbs/acre. In addition, the variability of RDM (as measured by standard errors in Figure 9a) is greatest in low quality habitat, suggesting that higher quality SKR habitat is characterized by consistently low RDM levels. In their analysis of these data, Spencer and Montgomery (2007) suggest that high quality SKR habitat is characterized by a threshold RDM level of less than 3,000 lbs/acre. Spring biomass and fall RDM were moderately correlated in 2006 (Figure 9b). Spring biomass was approximately twice the fall RDM levels in 2006.

#### *Photomonitoring data*

Representative photographs of the grasslands in the Preserve are provided in Attachment D (Figure D1). These photographs show the dominance of non-native grasses, primarily slender wild oat, in the Preserve. Note the taller grasses in the 2005 photograph indicative of the heavy rainfall of that year.

#### 4.1.3 Raptors

Detailed results of raptor surveys in the Preserve can be found in *Wintering Raptors of the Cagney Ranch and Surrounding Ramona Grasslands (2003-2006)* (WRI 2007, Appendix B), and are summarized in this section.

The grasslands and Santa Maria Creek riparian corridor in the Preserve support both wintering and breeding habitat for a diverse raptor community. A number of raptors detected in the Preserve are considered sensitive, including:

Bald eagle (*Haliaeetus leucocephalus*); SE, FP, County Group 1  
 Golden eagle (*Aquila chrysaetos*); FP, CSC, County Group 1  
 Ferruginous hawk; CSC, County Group 1  
 Northern harrier (*Circus cyaneus*); CSC, County Group 1  
 Merlin (*Falco columbarius*); CSC, County Group 1  
 Prairie falcon (*Falco mexicanus*); CSC, County Group 1  
 Barn owl (*Tyto alba*); County Group 2  
 Burrowing owl (*Athene cunicularia*); CSC, County Group 1  
 White-tailed kite (*Elanus leucurus*); FP, County Group 1  
 Cooper's hawk (*Accipiter cooperii*); CSC, County Group 1  
 Red-shouldered hawk (*Buteo lineatus*); County Group 1

(SE = California Endangered, FP = California Fully Protected, CSC = California Species of Concern)

A diverse assemblage of raptor species used the Preserve during winter months (Figure 10). WRI (2007) noted that the wintering data for 2005 were likely an under-estimate of the average abundance of raptors in the Preserve because data for February and March of that year were not

included in the average. The most abundant raptor species using the Preserve from 2003-2006, include American kestrel (*Falco sparverius*), ferruginous hawk, red-tailed hawk (*Buteo jamaicensis*), and turkey vulture (*Cathartes aura*). Burrowing owl, golden eagle, and prairie falcon also consistently used the Preserve, albeit in low numbers. Fewer than two burrowing owl individuals were typically seen during surveys from 2003-2005, but as many as five individuals were seen in 2006. Only one to two golden eagle and prairie falcon individuals were observed on any survey date. The maximum number ferruginous hawk observed in the Preserve from 2003-2006 was eight individuals (CBI 2007).

Raptors breeding in the Preserve included American kestrel, burrowing owl, barn owl, great-horned owl (*Bubo virginianus*), red-tailed hawk, red-shouldered hawk, and white-tailed kite (Figure 11). Red-shouldered hawk were by far the most frequent breeding species within and in areas immediately adjacent to the Preserve. These species generally used native trees along the Santa Maria Creek corridor and various non-native trees scattered within and around the Preserve (e.g., eucalyptus) for nesting, except for burrowing owls which used artificial nest boxes on the WRI property.

## 4.2 Vernal Pools, Vernal Swales, and Alkali Playas

### 4.2.1 Vernal pool, vernal swale, and alkali playa vegetation

Several rare plant species were detected in vernal pools and alkali playas in the Preserve (Figure 7), including:

Parish's brittlescale (*Atriplex parishii* var. *parishii*); County Group A

Coulter's saltbush (*Atriplex coulteri*); County Group A

Small-flower microseris (*Microseris douglasii* ssp. *platycarpha*); County Group D

#### *Species richness*

There was considerable variation in the plant species present in ephemeral aquatic habitats in the Preserve. Attachment F shows all species detected in each vernal pool surveyed in 2005 and 2006. A total of 98 taxa were recorded across all vernal pools; however, only 23 of these taxa (highlighted in Attachment F) were detected in more than half of the pools. Eleven of these 23 taxa are non-native.

Although there was substantial overlap, the vernal pool complexes considered in this report did show some unique species assemblages (Attachment F). For example, the reaches of the Cagney swale generally had higher taxonomic richness than other ephemeral aquatic habitats in the Preserve, including the highest number of species (42) recorded. Pools in the Airport complex were the only pools in the Preserve to support cryptantha (*Cryptantha* sp.), annual hairgrass (*Deschampsia danthonioides*), blue dicks (*Dichelostemma capitatum* ssp. *capitatum*), toothed downingia (*Downingia cuspidata*), smooth boisduvalia (*Epilobium pygmaeum*), dwarf peppergrass (*Lessingia filaginifolia* var. *filaginifolia*), grab lotus (*Lotus hamatus*), small-flower microseris (*Microseris douglasii* ssp. *platycarpha*), common muilla (*Muilla maritima*), dot-seed plantain (*Plantago erecta*), blue-eyed-grass (*Sisyrinchium bellum*), and tree clover (*Trifolium*

*ciliolatum*). The pools on Cagney Ranch (outside of the airport complex) were the only pools to support red maids (*Calandrinia ciliate*), American pillwort (*Pilularia americana*), and white tip clover (*Trifolium variegatum*). Although not all were detected in playa RAAP 100, the alkali playas were the only habitats to support Parish's brittlescale (*Atriplex parishii* var. *parishii*), Coulter's saltbush (*Atriplex coulteri*), and veiny peppergrass (*Lepidium oblongum* var. *insulare*).

#### *Species composition by pool zone*

The relative species composition varied both within pool zones and between the complexes (Figure 12). Across both years sampled, there was generally a lower percentage of bare ground within pool centers in the Airport complex relative to pools in the other complexes due to a high cover of Mediterranean barley (*Hordeum marinum* ssp. *gussoneanum*), annual beard grass (*Polypogon monspeliensis*), woolly marbles (*Psilocarphus brevissimus* var. *brevissimus*), long-beaked filaree, toothed downingia, and iris-leaved rush (*Juncus xiphiodes*). The pools in the Cumming swale had higher cover contributions from hairy clover fern (*Marsilea vestita*), and this species was only found in the center zone of pools where it was present. The cover of *Avena* and *Bromus* species were low in the centers of pools in all complexes.

Averaging across the two years sampled within the intermediate zone of the pools, species composition tended to be more uniform across complexes, although woolly marbles and iris-leaved rush did contribute more to the cover in the Airport complex than in other complexes. At the upland edges of the pools, the cover of slender wild oat and ripgut grass increased. Long-beaked filaree generally tended to increase in its contribution to cover moving from the center to upland edges of pools.

Total cover was generally higher in all complexes in 2005. The other notable change was the increase in cover of Bermuda grass (*Cynodon dactylon*) in the alkali playa (RAAP 100) in 2006 (Figure 12).

#### *Photomonitoring data*

Representative photographs of the vernal pools in the Preserve are provided in Attachment D (Figures D2). Figure D2 shows the conditions of vernal pool e44 in the Airport Complex. Note the extensive cover of non-native grasses.

#### 4.2.2 Hydrology and water characteristics

Detailed vernal pool hydrologic monitoring results are provided in *Biological Survey Report for the Ramona Grasslands Preserve* (RECON 2005, Appendix C). Vernal pool monitoring did not begin until well after the start of the rainy season in October 2004, and by January 21, 2005 nearly 17 inches of rain had been recorded at the Ramona Airport. Thus the trends summarized below are only representative of the latter half of the rainy 2004/2005 season.

Vernal pool hydrology varied by vernal pool complex (Figure 13), although the hydrology of individual pools within a complex was fairly variable. For example, the Cumming and Cagney swales supported the deepest and most persistent pools. One of the two pools sampled in the

Cumming swale (ev1) has been impounded, presumably to provide a water supply for cattle. The Cagney swale has been incised by runoff from the Ramona Airport tarmac, thus creating relatively deep and persistent pools. The monitored alkali playa (RAAP 100) is large relative to the other playas but also appears to sit within a swale that is fed by runoff from both the northern portion of Cagney Ranch and, as in 2005, periodic overbank flow from Santa Maria Creek. The vernal pools in the Cagney and Airport complexes were the shallowest and least persistent of all of the ephemeral aquatic habitats in the Preserve. The Airport complex has pools with “typical” vernal pool basin morphology (i.e., mima mound topography) and hydrologic function, whereas the Cagney pools mostly lie within the east-west trending swale (which also supports the alkali playas) and may have hydrology influenced by the network of swales in the Preserve (i.e., have a larger watershed area relative to the Airport complex) and by overbank flow from Santa Maria Creek. Vernal pools in the Airport complex dried earlier than pools in the other complexes; however, if pool e59, which does not lie in the east-west trending swale is removed from the analysis, the remaining three pools in the Cagney complex dried earlier than the Airport pools.

The conductivity of alkali playas measured in March 2005 ranged from 1,890 microsiemens ( $\mu\text{S}$ ) to over 1,990  $\mu\text{S}$ , whereas the conductivity of vernal pools in the Airport complex was less than 150  $\mu\text{S}$ . Water temperatures were quite variable during the 2005 survey period, ranging from 61 °F to 79 °F. Biweekly water temperature fluctuations of 4 °F to 6 °F within pools were common (RECON 2005, Appendix C). Dissolved oxygen concentrations of vernal pools were typically above 9 mg/l (RECON 2005, Appendix C), well above 100% saturation concentrations given the temperatures of the pools at the time of measurement. An anomalous reading of 2.4 mg/l was recorded for vernal pool e45 on April 1. Interestingly, the dissolved oxygen concentrations of the only alkali playa monitored (RAAP 100) ranged from 7.0 to 8.6 mg/l, uniformly lower than the dissolved oxygen concentrations measured in vernal pools on the same dates. Elevated concentrations of dissolved solids in water lower saturation concentrations, and the high conductivity of the alkali playas appears to be reflected in their lower dissolved oxygen concentrations.

#### 4.2.3 Fauna

Detailed results of faunal surveys are provided in *Biological Survey Report for the Ramona Grasslands Preserve* (RECON 2005, Appendix C). These results are summarized below for the two target taxa in this report, San Diego fairy shrimp and spadefoot toad. San Diego fairy shrimp is federally endangered and is a County Group 1 species. Western spadefoot toad is a California species of concern and a County Group 2 species.

##### *San Diego fairy shrimp*

San Diego fairy shrimp were commonly detected in the Preserve. While all ephemeral aquatic habitats had suitable ponding conditions for San Diego fairy shrimp (i.e., ponding for at least 2-weeks), San Diego fairy shrimp were only detected in pools in the Airport complex (pools e45, e46, e52, e53), the Cagney complex (pools e56, e58, e59, e62), one station in the Cagney swale (vs1), and the alkali playa (RAAP 100). No other fairy shrimp species were detected in the Preserve. Interestingly, while these pools retained water well into March or April, the last observation of fairy shrimp was on the February 4 survey date, and the majority of pools only

had fairy shrimp detected on the first survey date on January 21 (RECON 2005, Appendix C). Given the late start of vernal pool monitoring in the 2004/2005 rainy season, it is likely that many fairy shrimp may have already completed their life cycles prior to monitoring.

#### *Western spadefoot toad*

Western spadefoot toad larvae were detected in pools in the Airport complex (pools e46, e53, e77), the Cumming swale (ev1), and the Cagney swale (vs1, vs2, vs3, vs4). However, toadlets were only seen in pools e46 and vs3. It is not clear if spadefoot toads did not complete their lifecycles in the other pools where breeding was documented or if toadlets were just not detected at these pools. Two other amphibian species, western toad and Pacific tree frog, also successfully bred and metamorphosed in pools in the Cumming swale, Cagney vernal pools, Cagney swale, and the alkali playa (RAAP 100).

### 4.3 Santa Maria Creek Corridor

#### 4.3.1 Water quality

Complete water quality monitoring results for 2005-2006 are provided as Attachment G of this report. In this section, the baseline water quality monitoring results for Santa Maria Creek is summarized by groups of parameters. Water quality data are presented in three-dimensional graphs with date on the x-axis, the parameter values (e.g., concentration) on the y-axis, and a separate curve for each parameter at each station organized along the z-axis. For each parameter, stations are identified by a gradient of color from lightest (most upstream) to darkest (most downstream). Water quality monitoring stations are shown in Figure 5. Thus, the changes in parameter values across the different sampling dates can be compared side-by-side for each monitoring station.

#### *Temperature, dissolved oxygen, and pH*

Figure 14 shows the monitoring results for temperature, dissolved oxygen (DO), and pH for the three monitoring stations. Temperature of the creek changed in a similar fashion among the three stations across the monitoring period. As expected, temperatures generally increased over the course of the calendar year, with maximum temperatures seen in late spring. Two dips in temperature were seen in March and April of 2005, corresponding to storms that lowered temperatures. DO concentrations and pH exhibited similar patterns at all stations over all sampling dates.

#### *Stream flow and total suspended solids*

Stream flow and total suspended solids (TSS) are shown in Figure 15. Stream flow peaked in February 2005 following heavy winter rains and tapered off through the remainder of the year. Rainfall during the winter of 2006 was very low, and thus stream flow only exhibited a minimal increase during this year. TSS tends to correlate with discharge, as higher velocity flows can carry higher suspended sediment loads. TSS concentrations were highest at SMC1 and SMC2

(the most upstream and downstream stations), and these stations have much less vegetation cover in the channel than SMC3.

#### *Conductivity and total dissolved solids*

Figure 16 shows the monitoring results for conductivity and total dissolved solids (TDS). Since conductivity reflects the presence of dissolved ions, it is not surprising that TDS and conductivity are well correlated across the monitoring period, except for a dip in TDS in March of 2005. Both conductivity and TDS showed a spike at the end of 2005 at the SMC3 station.

#### *Total nitrogen and total phosphorus*

Total nitrogen (TN) and total phosphorus (P) are shown in Figure 17. These two elements show similar patterns across the monitoring period, but TN exhibits much higher variations in concentration. There also appears to be a gradient of TN through the project area, with higher TN upstream and lower downstream. This may reflect inputs from the urban and agricultural land uses upstream of the project area. The concentration of P shows peaks that are correlated with stream flow magnitude, as particulate P is often absorbed to sediment. Nutrient concentrations fall dramatically as stream flow declines during the late spring and summer of 2005.

#### *Total coliform bacteria*

Total coliform bacteria are shown Figure 18. Total coliform bacteria levels are highest at SMC 1, potentially an indication of agricultural runoff from areas upstream of the Ramona Grasslands. Bacterial levels tend to decline across all sampling stations during the late winter and spring, suggesting that bacteria are being flushed into the stream during rainfall events. The exception to this pattern is a peak in total coliform levels in late May 2005 at SMC1 (most upstream station), which is not related to increased stream flow.

#### *Chloride and sulfate*

The concentrations of chloride and sulfate ions are shown in Figure 19. These ions exhibit very low concentrations early in 2006 but steadily increase over the later winter and spring of this year. The highest concentrations of chloride were seen at station SMC1 during later 2005 and 2006, possibly reflecting agricultural runoff.

### 4.3.2 Channel geomorphology

Cross-sections graphically displaying the geomorphology of the Santa Maria Creek channel are provided in Figures 20a and 20b. The Santa Maria Creek channel within the Preserve varies in width from 30 to 200 ft and in depth from 3 to 12 ft, although the active channel is generally confined to a small portion of the wider reaches of the channel (e.g., SG9, SG10, and SG11, Figure 20b). The channel generally becomes wider and more incised at the western end of the preserve.

#### 4.3.3 Riparian vegetation

Riparian tree species in the Santa Maria Creek corridor largely consist of red willow (*Salix laevigata*), arroyo willow (*Salix lasiolepis*), black willow (*Salix gooddingii*), and Fremont cottonwood (*Populus fremontii*), with coast live oak (*Quercus agrifolia*) and western sycamore (*Platanus racemosa*) present at the western end of the Preserve. Riparian shrub species are dominated by mulefat (*Baccharis salicifolia*), and young willows, including narrow-leaf willow (*Salix exigua*) at the eastern end of the Preserve. The herbaceous layer at the western end of the Preserve is dominated by freshwater marsh species, including Olney's bulrush (*Schoenoplectus americanus*), yerba mansa (*Anemopsis californica*), Mexican rush (*Juncus mexicanum*), western ragweed (*Ambrosia psilostachya*), while upstream of Rangeland Road herbaceous species are dominated by yerba mansa, Mexican rush, mugwort (*Artemisia douglasiana*), water speedwell (*Veronica anagallis-aquatica*), cocklebur (*Xanthium strumarium*), great marsh evening primrose (*Oenothera elata* ssp. *hirsutissima*), creeping wild rye (*Leymus triticoides*), grass poly (*Lythrum hyssopifolium*), white sweet clover (*Melilotus alba*) and mulefat seedlings.

The Santa Maria Creek corridor currently has low vegetative cover, particularly tree cover (Figures 21a, b). Riparian transects 3, 4, and 5 have the highest tree cover in the Preserve; although the cover recorded at Transect 3 is not indicative of that stream reach, which actually has relatively little tree cover. Shrub cover is more uniform at 10-25%, except for Transects 2, 3, and 8 that have no shrub cover. Riparian tree and shrub species were producing seedlings each year (Table 7); however, the survival rate of these seedlings is unknown.

**Table 7. Total number of seedlings recorded at each riparian transect location.**

<b>Transect</b>	<b><i>Mulefat</i></b>	<b><i>Coast live oak</i></b>	<b><i>Black willow</i></b>	<b><i>Red willow</i></b>	<b><i>Arroyo willow</i></b>	<b>Transect Total</b>
<b>1</b>	87					87
<b>2</b>	9					9
<b>3</b>			1			1
<b>4</b>	84					84
<b>5</b>	527	1				528
<b>6</b>	303		7			310
<b>7</b>	110			1	3	114
<b>8</b>						
<b>9</b>						
<b>10</b>	10		3			13
<b>11</b>	41		1		0	42
<b>Species Total</b>	<b>1,171</b>	<b>1</b>	<b>12</b>	<b>1</b>	<b>3</b>	<b>1,188</b>

#### *Photomonitoring data*

Representative photographs of the riparian habitats in the Preserve are provided in Attachment D (Figures D3-D5). Figure E3 shows the virtual absence of riparian vegetation in Reach E of Santa Maria Creek as a result of historic cattle grazing. Figure D4 shows the woodland and shrubby

understory structure of the riparian vegetation in Reach D of Santa Maria Creek. Figure D5 shows the mulefat scrub and freshwater marsh communities in Reach A of Santa Maria Creek.

#### 4.3.4 Avifauna

Detailed avifauna survey results are provided in *Biological Survey Report for the Santa Maria Creek Restoration Project: riparian birds* (Lovio 2007, Appendix D) and are summarized here.

Several sensitive species were detected in riparian habitats in the Preserve, including:

Loggerhead shrike (*Lanius ludovicianus*); CSC, County Group 1

Southwestern willow flycatcher (*Empidonax traillii estimus*)\*; FE, SE, County Group 1

Western bluebird (*Sialia mexicana*); County Group 2

Yellow warbler (*Dendroica petechia*); CSC, Group 2

(FE = Federal Endangered, SE = California Endangered, CSC = California Species of Concern)

\* Note: A single, singing willow flycatcher was detected in the Preserve on a single date in June, but did not establish a breeding territory and was considered to be migrating through the Preserve. However, this species is considered to potentially occur in the Preserve.

Bird species most commonly breeding within the Santa Maria Creek corridor are primarily habitat generalists, and include red-tailed hawk, American kestrel, mourning dove (*Zenaidura macroura*), Anna's hummingbird (*Calypte anna*), black phoebe (*Sayornis nigricans*), ash-throated flycatcher (*Myiarchus cinerascens*), Cassin's kingbird (*Tyrannus vociferans*), western kingbird (*Tyrannus verticalis*), western scrub-jay (*Aphelocoma californica*), American crow (*Corvus brachyrhynchos*), bushtit (*Psaltiriparus minimus*), spotted towhee (*Pipilo maculatus*), California towhee (*Pipilo crissalis*), brown-headed cowbird (*Molothrus ater*), blue grosbeak (*Guiraca caerulea*), and house finch (*Carpodacus mexicanus*). Only one species detected in 2005, the yellow warbler (*Dendroica petechia*), may be regarded as a riparian obligate species. However, this species can show affinity for non-native forests, and was associated with the eucalyptus woodland adjacent to the creek (Lovio 2006). In addition, several species, including sora (*Porzana carolina*), Virginia rail (*Rallus limicola*), and red-winged blackbird (*Agelaius phoeniceus*), were associated with the mulefat scrub and marsh habitats in the westernmost reach of the creek corridor.

#### 4.3.5 Arroyo toad

Detailed arroyo toad survey results are provided in *Biological Survey Report for the Ramona Grasslands Preserve* (RECON 2005, Appendix C) and *Biological Survey/Monitoring Report for the Santa Maria Creek Restoration Project: arroyo toads* (Hollingsworth et al. 2006, Appendix E) and are summarized here.

A breeding population of arroyo toad has been documented in the Preserve, but only in the western reach of Santa Maria Creek, downstream of Rangeland Road (Figure 22). Arroyo toads were detected in this reach in both 2005 and 2006 (RECON 2005 and Hollingsworth 2006, Appendices C and E), but their abundance was considered to be less than is typical for similar stream systems (Hollingsworth et al. 2006, Appendix E). Western toad, two-striped garter snake,

California kingsnake (*Lampropeltis getula*), and Pacific treefrog were also detected in this stream reach. A number of non-native species including crayfish (*Procambarus clarkii*), mosquitofish (*Gambusia affinis*), largemouth bass (*Micropterus salmoides*), green sunfish (*Lepomis cyanellus*), and bullfrog (*Rana catesbeiana*) are present in the reach of the creek occupied by arroyo toads. The abundance of bullfrogs was considered quite high (Hollingsworth et al. 2006, Appendix E).

#### 4.4 Invasive Non-native Plant Removal

Detailed results of ongoing invasive non-native plant control efforts are provided in *Biological Survey Report for the Ramona Grasslands Preserve* (RECON 2005, Appendix C) and *Invasive Weed Report for the Santa Maria Creek Restoration Project: grassland and riparian invasive weed control efforts and results* (Kelly & Associates 2007, Appendix F), and are summarized below.

The locations of invasive non-native plants within the Preserve are shown in Figure 23a-c. Of particular concern were artichoke thistle (*Cynara cardunculus*), salt cedar (*Tamarix ramosissima*), and giant reed (*Arundo donax*). Additional invasive non-native plants, detected during the removal efforts in 2005 and 2006, include milk thistle (*Silybum marianum*), Italian thistle (*Carduus pycnocephalus*), intermediate wheatgrass (*Elytrigia intermedia*), perennial pepperweed (*Lepidium latifolium*), and minor amounts of other non-native species (Figure 23a, b). The majority of these invasive, non-native plants have been treated in 2005 and 2006 (Figure 24), and additional treatment will occur at least into the winter of 2007.

#### 5.0 DISCUSSION

##### *Preserve-wide*

The approximately 2,000-acre Ramona Grasslands Preserve supports a wide diversity of plant and animals species, including a number of rare, threatened, or endangered species (Attachments B and C). The Preserve provides regionally significant grassland habitat for animal species such as SKR, ferruginous hawk, and golden eagle, and native plant species such as purple needlegrass, San Diego thornmint, small-flowered morning glory, and round-leaved filaree. San Diego fairy shrimp and spadefoot toad are well-distributed in ephemeral aquatic habitats (i.e., vernal pools, vernal swales, and alkali playas) in the Preserve. Santa Maria Creek supports a population of arroyo toad and both marsh and riparian associated bird species. Unique alkali playas are present in the Preserve, which provide habitat for rare plants species such as Parish's brittle scale and Coulter's saltbush.

##### *Grasslands*

The distribution of sensitive grassland species in the Preserve appears to be related to the structure and composition of grassland habitats, which a function of soil composition, topography, and grazing pressure. For example, SKR habitat is largely located within the central and northern portions of the Preserve (Figure 6), which are characterized by well-drained, loamy soil types (Figure 3) and heavy cattle grazing. Grassland habitats in these areas tend to have higher amounts of bare ground and greater ratios of forb species to grass species (Spencer and Montgomery 2007, Appendix A). These areas are also prime raptor foraging habitat, as the

greater amount of bare ground and forb cover provide a habitat structure more conducive to detecting and catching prey species, primarily small mammals. The abundance of SKR is also sensitive to plant biomass, with areas of higher biomass or RDM exhibiting lower densities of SKR (Figure 9a, Appendix A). SKR abundance within a given year is dynamic in response to fluctuations in grassland structure, and the distribution of SKR shrinks back to the highest quality habitat areas in years of higher rainfall and increased vegetative growth.

Grasslands on clayey soil types, often located around the margins of the Preserve and presumably receiving less grazing pressure, are not good habitat for SKR. These clayey grassland habitats support the majority of the rare native plant species in the Preserve, and exhibit higher plant cover, less bare ground, and greater RDM than do areas supporting high densities of SKR. Thus, distinguishing between loamy and clayey grassland habitats appears to be useful means of identifying management targets and desired conditions for the grasslands in the Preserve. In addition, SKR habitat quality can be characterized by metrics of grassland structure. For example, high quality SKR habitat was characterized by mean RDM of 1,800 lbs/acre, and Spencer and Montgomery (2007, Appendix A) suggest that RDM of less than 3,000 lbs/acres may be a threshold for suitable SKR habitat. In addition, positive correlations between spring grassland biomass and fall and winter RDM were observed in this study (Figure 9b); thus, spring biomass measurements may provide land managers with early information to adjust grazing intensity to achieve management objectives.

The Preserve provides important habitat for wintering and breeding raptors. Ferruginous hawks are regular visitors to the Preserve, with between four to eight individuals regularly observed from 2003-2006; although WRI estimated as many as 22 ferruginous hawks were using the preserve in 2005 (WRI 2007, Appendix B). A pair of golden eagles regularly forages in the Preserve, and presumably these are the eagles nesting in nearby Bandy Canyon. Red-shouldered hawks were the most frequent breeders in the Preserve, and they and other species make use of the native and non-native trees in the Preserve for nesting. Burrowing owls were reintroduced into the Preserve by WRI in 2005 and successfully breed in artificial nest boxes that year (WRI 2007, Appendix B). The average number of burrowing owls observed in 2006 was significantly greater than in previous years, indicating that the reintroduced birds and their offspring are likely over-wintering in the Preserve.

#### *Vernal pools, vernal swales, and alkali playays*

The various classes of ephemeral aquatic habitats in the Preserve exhibit unique biological structures and functions. For example, the Cagney swale supported the highest number of plant species but at least a dozen native plant species occurred in the Airport vernal pool complex that were detected in no other ephemeral aquatic habitats in the Preserve. The center and intermediate zones of the Airport pools, which support the majority of the native vernal pool flora, also had the greatest total plant cover and thatch cover (Figure 12). The alkali playays in the Preserve were the only habitat supporting Parish's brittlescale and Coulter's saltbush, and are also distinctive from other ephemeral aquatic habitats in their species composition and water chemistry. The hydrology of the various classes of ephemeral aquatic habitats also varied substantially, with pools associated with swales exhibiting much longer periods of inundation than the mima mound vernal pools in the Airport complex (Figure 13).

San Diego fairy shrimp and western spadefoot toads were well-distributed in the ephemeral aquatic habitats in the Preserve. Interestingly, San Diego fairy shrimp were only observed for a short period of time at the beginning of the survey period in 2005. Since vernal pool monitoring in 2005 started after significant fall season rains in 2004, it is unknown when San Diego fairy shrimp first emerged in vernal pools in the Preserve. Fairy shrimp were detected in all types of ephemeral aquatic habitats in the Preserve; although it is unknown whether fairy shrimp were transported into the single alkali playa monitored during 2005 by overland flow in the extremely wet winter that year or naturally occur in alkali playas. In contrast, western spadefoot toad was not detected in alkali playas, but Pacific treefrog and western toad were.

#### *Santa Maria Creek corridor*

Water quality data collected in 2005 and 2006 suggest that runoff from upstream urban and agricultural areas may be a non-point source of some pollutants in Santa Maria Creek. For example, conductivity, TDS, TN, total coliform bacteria, and chloride all exhibited their highest levels at SMC1, the most upstream monitoring station. The baseline water quality monitoring also showed that the concentrations of a number of water quality parameters, including TN, P, and total coliform bacteria, are correlated with stream flow, indicating that they may be transported into the system from upstream areas. The concentration of TSS is also correlated with stream flow but is lowest at the central sampling station, which is located in the most vegetated portion of Santa Maria Creek within the project area. This result tends to support the restoration project's objective of improving water quality by restoring riparian vegetation and riverine functions. It is anticipated that improvements in water quality in Santa Maria Creek will be realized once future land management actions are implemented, and that the water quality monitoring program and baseline data presented here will provide a means for assessing these improvements.

Riparian vegetation in the Preserve is severely degraded by years of cattle grazing. Currently, there is virtually no tree cover and little shrub cover in the majority of reaches of Santa Maria Creek (Figure 21a). Only in the reach traversing the private properties along Vorhees Lane, where cattle have been excluded since the 1970s (CBI 2006), has a riparian woodland structure established. The cottonwood-willow riparian woodland in this reach is considered to be the model of what riparian habitat in Santa Maria Creek upstream of Rangeland Road should look like in the future following elimination of cattle from the creek corridor. As of the end of 2006, almost the entire Santa Maria Creek corridor has been fenced to exclude cattle. Seedling riparian shrub and tree species, principally the understory shrub mulefat but also overstory willow species, were observed within transects surveyed for this report. Presumably seedling survival rate and vegetative cover will increase in the future. In addition, the Santa Maria Creek corridor in the western end of the Preserve supports perennial surface water and marsh plant communities that are distinct from the willow and mulefat dominated plant communities in the upstream reaches of the Creek exhibiting ephemeral surface water flow.

Riparian birds currently using the Preserve are dominated by generalist species. Only a single obligate riparian species was observed (yellow warbler), which was found nesting in non-native forest adjacent to the creek. Enhancement of habitats in the Preserve for riparian obligate bird species will likely not result in the loss of habitat generalists from the avian community, but rather in an increase in bird species diversity by the addition of obligate riparian species. In

addition, a distinct avifauna associated with the marsh habitats in the Santa Maria Creek corridor downstream of Rangeland Road was documented (Lovio 2007, Appendix D). The habitat for these marsh-associated species, which include sora, Virginia rail, and red-winged black bird, should be considered a management target distinct from the willow and mulefat dominated habitats and their associated riparian avifauna that occur upstream of Rangeland Road

Arroyo toad habitat in the Preserve appears to be restricted to the reach of Santa Maria Creek downstream of Rangeland Road, and is driven by surface water hydrology. The creek channel in the upstream portion of the Preserve is predominately sandy alluvium and is characterized by an ephemeral stream flow. In the reaches upstream of Rangeland Road, Santa Maria Creek flows only in response to rainfall events. Even in the very wet winter of 2005, surface flow had stopped by June. However, at the extreme western end of the Preserve, surface water appears to be perennial. CBI (2006) speculated that the granitic geology forming the western “wall” of the Santa Maria Valley, through which the creek has cut Bandy Canyon, may maintain groundwater at a shallower depth below the channel bottom than in the eastern end of the Valley by “impounding” shallow groundwater as it flows through the Valley to Bandy Canyon. This would also explain the presence of marsh vegetation in this reach of the creek, which requires consistently high groundwater. Regardless of the mechanism, which clearly requires more investigation, it appears that the western end of the Preserve consistently provides suitable habitat for arroyo toads, whereas the eastern end of the Preserve typically does not provide suitable habitat for this species. In addition, Hollingsworth et al. noted a very high abundance of bullfrogs and other non-native aquatic species in the arroyo toad habitat in the Preserve and observed that arroyo toad abundance was less than is typical for similar stream systems occupied by this species (Hollingsworth et al. 2006, Appendix E). The Ramona Water District operates effluent holding ponds in close proximity to arroyo toad habitat in Santa Maria Creek that could be a source of bullfrogs. Nonnative predators in Santa Maria Creek, such as bullfrogs, may be adversely affecting the abundance of arroyo toads in this system.

#### *Invasive non-native plants*

Invasive non-native plants, particularly artichoke thistle, continue to be a problem in the Preserve. Invasive, non-native plants have been the subject of eradication efforts since 2005, and as of the writing of this report, artichoke thistle continues to germinate in significant numbers. Ongoing monitoring and control of invasive non-native plants will be required for at least several years to adequately control these species. In addition, adjacent properties outside of the Preserve continue to serve as sources of invasive non-native plants, and every effort should be made to expand weed abatement efforts to these areas.

## 6.0 REFERENCES

- Bailey, R.G., P.E. Avers, T. King, and W.H. McNab (eds.). 1994. Ecoregions and subregions of the United States. Color map at 1:750,000, accompanied by a supplementary table of map unit descriptions. Prepared for USDA, Forest Service. U.S. Geological Survey, Washington, DC.
- Bibby, C.J., N.D. Burgess, & D.A. Hill. 1992. Bird Census Techniques. Academic Press, London.

- City of San Diego Water Department. 2004. Santa Maria Creek Restoration Water Monitoring Quality Assurance Project Plan
- Conservation Biology Institute (CBI) 2006. Santa Maria Creek riparian habitat and channel geomorphology assessment. Prepared for The Nature Conservancy. June.
- Conservation Biology Institute (CBI) 2007. Baseline conditions report for the Ramona Grasslands Preserve. Prepared for the County of San Diego Department of Parks and Recreation. January. Volume 2. Technical Appendices.
- Federal Aviation Administration (FAA). 2002. Habitat management plan (HMP) for the Ramona Airport property: Stephens' kangaroo rat. November. 64 pp + appendix.
- Federal Aviation Administration (FAA). 2003. Ramona Airport Improvement Project Vernal Pool Habitat Management Plan. October. 64pp + appendices.
- Goudey, C.B., and D.W. Smith (eds.). 1994. Ecological units of California: subsections. Color map at 1:1,000,000. USDA, Forest Service, San Francisco, CA.
- Haas, W.E. and M.J. O'Farrell. 2005. Final Report for the 2005 Stephens' Kangaroo Rat Monitoring Program for the Ramona Airport Improvement Project. December 2005. Prepared for County of San Diego Department of Public Works, Environmental Services Unit, San Diego, CA. vi+29pp+appendices.
- Hollingsworth, B.D., C.M.Shaffer, and M. Roll. 2006. Biological survey report for the Santa Maria Creek Restoration Project: arroyo toads. Prepared for County of San Diego Department of Parks and Recreation. December.
- Kelly & Associates. 2007. Invasive weed report for the Santa Maria Creek Restoration Project: grassland and riparian invasive weed control efforts and results. Prepared for County of San Diego Department of Parks and Recreation. January.
- Lovio, J.C. 2007. *Biological Survey Report for the Santa Maria Creek Restoration Project: riparian birds*. Prepared for County of San Diego Department of Parks and Recreation. January.
- McFedries, M. 2006. Personal communication to Jennifer Haines.
- McNab, W.H., and P.E. Avers (eds.). 1994. Ecological subregions of the United States: section descriptions. USDA, Forest Service Publication WO-WSA-5. Washington, DC.
- Miles, S.R., and C.B. Goudey. 1998. Ecological subregions of California: section and subsection descriptions. USDA Forest Service, Pacific Southwest Region. R5-EM-TP-005-NET.
- O'Farrell, M.J. 1992. Establishment of a population monitoring program for the endangered Stephens' kangaroo rat. Trans. Western Section Wildlife Society 28:112-119.
- O'Farrell, M.J. 2000a. Untitled report concerning surveys for Stephens' kangaroo rat on the Cumming Ranch Special Planning Area. Prepared for Ecoventures California. 28 August 2000.
- O'Farrell, M.J. 2000b. Untitled report concerning surveys for Stephens' kangaroo rat on the Highland Valley Estates. Prepared for Mooney & Associates. 21 November 2000.

- O'Farrell, M.J. 2002. Untitled report concerning surveys for Stephens' kangaroo rat on the Highland Valley Estates. Prepared for Mooney & Associates. 26 January 2002.
- O'Farrell, M.J. 2004. Untitled report concerning surveys for Stephens' kangaroo rat on the Cumming Ranch Special Planning Area. Prepared for Ecoventures California. 21 February 2004.
- Ogden Environmental and Energy Services. 1998. Stephens' kangaroo rat study for the Ramona Airport expansion project, Ramona, California. Prepared for KEA Environmental and County of San Diego Department of Public Works.
- Price, M.V., W.S. Longland, and R.L. Goldingay. 1991. Niche relationships of *Dipodomys agilis* and *D. stephensi*: Two sympatric kangaroo rats of similar size. *American Midland Naturalist* 126:172-186.
- Ralph, C.J., G.R. Geupel, P.Pyle, T.E. Martin, & D.F. DeSante. 1993. Handbook of field methods for monitoring landbirds. Gen. Tech. Report PSW-GTR—144. Albany, CA: Pacific Southwest Research Station, Forest Service, U.S. Department of Agriculture.
- RECON. 2005. Biological survey report for the Ramona Grasslands Preserve in Ramona, California. Prepared for County of San Diego Department of Parks and Recreation. October.
- Spencer, W.D. 2002. Final habitat management plan (HMP) for the Ramona Airport property: Stephens' kangaroo rat. Prepared for U.S. Department of Transportation, Federal Aviation Administration, and County of San Diego Department of Public Works. November.
- Spencer, W.D and S.J. Montgomery. 2007. Biological Survey Report for the Santa Maria Creek Restoration Project: Stephens' kangaroo rat. Prepared for County of San Diego Department of Parks and Recreation. January.
- Tellam, S. 2004. Personal Communication to Michael White
- U.S. Department of Agriculture (USDA), Soil Conservation Service and Forest Service. 1973. Soil survey San Diego area, California.
- 1996 Interim Survey Guidelines to Permittees for Recovery Permits under Section 10(a)(1)(A) of the Endangered Species Act for the Listed Vernal Pool Branchiopods.
- U.S. Fish and Wildlife Service. 1999. Survey Protocol for the Arroyo Toad.
- U.S. Geological Survey (USGS). 2005. Monitoring Protocol for Arroyo Toads (*Bufo californicus*). Sara L. Compton and Robert N. Fisher. USGS Western Ecological Research Center. Draft Field Protocol. March.
- Ecological Research Center, Draft Field Protocol. March 2005. Van Velzen, W.T. 1972. Breeding-bird census instructions. *American Birds* 26(6):927-931.
- Western Regional Climate Center. 2006. Southern California climate summaries. <http://www.wrcc.dri.edu/summary/climsmsca.html>
- Wildlife Research Institute (WRI). 2007. Wintering raptors of the Cagney Ranch and surrounding Ramona Grasslands (2003-2006). Prepared for County of San Diego Department of Parks and Recreation. January.




Figure 1. Project Location.




Figure 2. Conserved Properties.




Figure 3. Soil series within the Preserve.




Figure 4a. Location of Preserve within the San Dieguito Watershed.




Figure 4b. Estimated extent of the flooding of Santa Maria Creek in 2005.




Figure 5. Locations of field monitoring stations in the Preserve. Note: there are no grassland monitoring plots 24-29.




Figure 6. Density of SKR burrows and delineation of core SKR management areas in the Preserve.




Figure 7. Locations of rare plants and patches of purple needlegrass in the Preserve.




Figure 8. Percent cover of grassland species within categories of SKR habitat quality (i.e., low medium, and high). Note: thatch cover data are missing for 2005.




Figure 9a. Average biomass ( $\pm$  SE) by SKR habitat quality.




Figure 9b. Spring and fall 2006 biomass correlation.




Figure 10. Average abundance of wintering raptors in the Preserve.




Figure 11. Raptor species breeding in the Preserve.




Figure 12. Vernal pool cover by pool zone in 2005 and 2006. Note: thatch cover data are missing for 2005.




Figure 13. Average depth of vernal pools by complex in 2005.




Figure 14. Temperature, dissolved oxygen, and pH at monitoring stations SMC1, SMC2, and SMC3 for the 2005-2006 monitoring period.




Figure 15. Stream flow and total suspended solids at monitoring stations SMC1, SMC2, and SMC3 for the 2005-2006 monitoring period.




Figure 16. Total dissolved solids and conductivity at monitoring stations SMC1, SMC2, and SMC3 for the 2005-2006 monitoring period.




Figure 17. Total nitrogen and total phosphorus at monitoring stations SMC1, SMC2, and SMC3 for the 2005-2006 monitoring period.




Figure 18. Total coliform bacteria at monitoring stations SMC1, SMC2, and SMC3 for the 2005-2006 monitoring period.




Figure 19. Chloride and sulfate at monitoring stations SMC1, SMC2, and SMC3 for the 2005-2006 monitoring period.




Figure 20a. Channel geomorphology at cross-section SG2-SG6.




Figure 20b. Channel geomorphology at cross-section SG7-SG11.




Figure 21a. Tree and shrub cover at riparian transect monitoring locations in 2005.




Figure 21b. Tree and shrub cover at riparian transect monitoring locations in 2006.




Figure 22. Distribution of arroyo toads detected in the Preserve.




Figure 23a. Locations of invasive non-native plant species in the Preserve.




Figure 23 b. Locations of invasive non-native plant species in the Preserve.




Figure 24. Aggressive non-native species treated in the Preserve.

## **ATTACHMENTS**

ATTACHMENT A—PLANT SPECIES LIST

ATTACHMENT B—ANIMAL SPECIES LIST

ATTACHMENT C—PHOTOGRAPHS OF SELECTED SENSITIVE SPECIES

ATTACHMENT D—PHOTOMONITORING RESULTS

ATTACHMENT E—GRASSLAND SPECIES RICHNESS

ATTACHMENT F—VERNAL POOL SPECIES RICHNESS

ATTACHMENT G—WATER QUALITY RESULTS

# **ATTACHMENT A**

## **PLANT SPECIES LIST**

## ATTACHMENT A PLANT SPECIES LIST

Common Name	Scientific Name	Federal	State	County of San Diego
San Diego thornmint	<i>Acanthomintha ilicifolia</i>	FT	SE	A
Blow-wives	<i>Achyrachaena mollis</i>			
Desert crested wheatgrass	<i>Agropyron desertorum</i>			
Pigweed	<i>Amaranthus</i> sp.			
Western ragweed	<i>Ambrosia psilostachya</i>			
Fiddleneck	<i>Amsinckia</i> sp.			
Scarlet pimpernel	<i>Anagallis arvensis</i>			
Yerba mansa	<i>Anemopsis californica</i>			
Mayweed	<i>Anthemis cotula</i>			
Nuttall's snapdragon	<i>Antirrhinum nuttallianum</i> ssp. ?			
Common celery	<i>Apium graveolens</i>			
California sagebrush	<i>Artemisia californica</i>			
Mugwort	<i>Artemisia douglasiana</i>			
Tarragon	<i>Artemisia drancunculus</i>			
Giant reed	<i>Arundo donax</i>			
California milkweed	<i>Asclepias californica</i>			
Coulter's saltbush	<i>Atriplex coulteri</i>			A
Parish's brittlescale	<i>Atriplex parishii</i> var. <i>parishii</i>			A
Slender wild oat	<i>Avena barbata</i>			
Wild oat	<i>Avena fatua</i>			
Mule fat	<i>Baccharis salicifolia</i>			
Black mustard	<i>Brassica nigra</i>			
California brickellbush	<i>Brickellia californica</i>			
Dwarf brodiaea	<i>Brodiaea terrestris</i> ssp. <i>kernensis</i>			
Ripgut grass	<i>Bromus diandrus</i>			
Soft chess	<i>Bromus hordeaceus</i>			
Foxtail chess	<i>Bromus madritensis</i> ssp. <i>rubens</i>			
Red maids	<i>Calandrinia ciliata</i>			
California large-leaf filaree	<i>California macrophylla</i>			B
Round-leaved filaree	<i>California macrophylla</i>			B
Water-starwort	<i>Callitriche</i> sp.			
Splendid mariposa lily	<i>Calochortus splendens</i>			
Morning-glory	<i>Calystegia macrostegia</i> ssp. ?			
Primrose	<i>Camissonia claviformis</i> ssp. ?			
Sandysoil sun cup	<i>Camissonia strigulosa</i>			
Rusty sedge	<i>Carex subfusca</i>			
Safflower	<i>Carthamus tinctorius</i>			
Owl's-clover	<i>Castilleja densiflora</i> ssp. <i>densiflora</i>			
Purple owl's-clover	<i>Castilleja exserta</i> ssp. <i>exserta</i>			
Tocalote	<i>Centaurea melitensis</i>			
Canchalagua	<i>Centaureum venustum</i>			

Common Name	Scientific Name	Federal	State	County of San Diego
Southern tarplant	<i>Centromadia parryi</i> ssp. <i>australis</i>			A
Common chaffweed	<i>Centunculus minimus</i>			
Mouse-ear chickweed	<i>Cerastium glomeratum</i>			
Spotted spurge	<i>Chamaesyce maculata</i>			
Pigweed	<i>Chenopodium</i> sp.			
Soap plant	<i>Chlorogalum parviflorum</i>			
Mediterranean chicory	<i>Cichorium intybus</i>			
Clarkia	<i>Clarkia purpurea</i> ssp. ?			
Miner's-lettuce	<i>Claytonia perfoliata</i> ssp. ?			
Virgin's bower	<i>Clematis</i> sp.			
Small-flower bindweed	<i>Convolvulus simulans</i>			D
Horseweed	<i>Conyza</i> sp.			
Common sand-aster	<i>Corethrogyne filaginifolia</i> var. <i>filaginifolia</i>			
California-aster	<i>Corethrogyne filaginifolia</i> var. <i>filaginifolia</i>			
African brass-buttons	<i>Cotula coronopifolia</i>			
Water pygmyweed	<i>Crassula aquatica</i>			
Pygmyweed	<i>Crassula connata</i>			
Alkali weed	<i>Cressa truxillensis</i>			
Dove weed	<i>Croton setigerus</i>			
Prickle grass	<i>Crypsis schoenoides</i>			
Cryptantha	<i>Cryptantha</i> sp.			
Coyote melon	<i>Cucurbita palmata</i>			
Dodder	<i>Cuscuta</i> sp.			
Artichoke thistle	<i>Cynara cardunculus</i>			
Bermuda grass	<i>Cynodon dactylon</i>			
Nutsedge	<i>Cyperus</i> sp.			
Western jimson weed	<i>Datura wrightii</i>			
Rattlesnake weed	<i>Daucus pusillus</i>			
Fascicled tarweed	<i>Deinandra fasciculata</i>			
Annual hairgrass	<i>Deschampsia danthonioides</i>			
Blue dicks	<i>Dichelostemma capitatum</i> ssp. <i>capitatum</i>			
Saltgrass	<i>Distichlis spicata</i>			
Padre's shooting star	<i>Dodecatheon clevelandii</i> ssp. <i>clevelandii</i>			
Toothed downingia	<i>Downingia cuspidata</i>			
Mexican tea	<i>Dysphania ambrosioides</i>			
Pale spike-rush	<i>Eleocharis macrostachya</i>			
Tall wheatgrass	<i>Elytrigia pontica</i> ssp. <i>pontica</i>			
Willow herb	<i>Epilobium ciliatum</i> ssp. <i>ciliatum</i>			
Smooth boisduvalia	<i>Epilobium pygmaeum</i>			
Tall buckwheat	<i>Eriogonum elongatum</i> var. <i>elongatum</i>			

Common Name	Scientific Name	Federal	State	County of San Diego
Coast California buckwheat	<i>Eriogonum fasciculatum</i> var. <i>fasciculatum</i>			
Mountain buckwheat	<i>Eriogonum fasciculatum</i> var. <i>polifolium</i>			
Long-beak filaree	<i>Erodium botrys</i>			
Red-stem filaree	<i>Erodium cicutarium</i>			
White-stem filaree	<i>Erodium moschatum</i>			
Reticulate-seed spurge	<i>Euphorbia spathulata</i>			
Narrow-leaf filago	<i>Filago gallica</i>			
Bedstraw	<i>Galium angustifolium</i>			
Nit grass	<i>Gastridium ventricosum</i>			
Carolina geranium	<i>Geranium carolinianum</i>			
Purple-spot gilia	<i>Gilia clivorum</i>			
California everlasting	<i>Gnaphalium californicum</i>			
Rayless gumplant	<i>Grindelia camporum</i> var. <i>bracteosa</i>			
Barbgrass	<i>Hainardia cylindrica</i>			
Saw-toothed goldenbush	<i>Hazardia squarrosa</i> var. <i>grindelioides</i>			
Crete hedypnois	<i>Hedypnois cretica</i>			
Peak rush-rose	<i>Helianthemum scoparium</i>			
Western sunflower	<i>Helianthus annuus</i>			
Salt heliotrope	<i>Heliotropium curassavicum</i>			
Telegraph weed	<i>Heterotheca grandiflora</i>			
Goldenaster	<i>Heterotheca sessiliflora</i>			
Short-pod mustard	<i>Hirschfeldia incana</i>			
Graceful tarplant	<i>Holocarpha virgata</i> ssp. <i>elongata</i>			D
Little barley	<i>Hordeum intercedens</i>			C
Mediterranean barley	<i>Hordeum marinum</i> ssp. <i>gussoneanum</i>			
Hare barley	<i>Hordeum murinum</i> ssp. <i>leporinum</i>			
Smooth cat's-ear	<i>Hypochaeris glabra</i>			
Spreading goldenbush	<i>Isocoma menziesii</i> var. <i>menziesii</i>			
Howell's quillwort	<i>Isoetes howellii</i>			
Mexican rush	<i>Juncus arcticus</i> var. <i>mexicanus</i>			
Toad rush	<i>Juncus bufonius</i> var. <i>bufonius</i>			
Mariposa rush	<i>Juncus dubius</i>			
Iris-leaved rush	<i>Juncus xiphiodes</i>			
Prickly lettuce	<i>Lactuca serriola</i>			
Goldentop	<i>Lamarckia aurea</i>			
Dwarf peppergrass	<i>Lepidium latipes</i> var. <i>latipes</i>			
Shining peppergrass	<i>Lepidium nitidum</i> var. <i>nitidum</i>			
Veiny peppergrass	<i>Lepidium oblongum</i> var. <i>insulare</i>			
Creeping wild rye	<i>Leymus triticoides</i>			
Flowering-quillwort	<i>Lilaea scilloides</i>			

Common Name	Scientific Name	Federal	State	County of San Diego
Southern mudwort	<i>Limosella acaulis</i>			
Blue toad flax	<i>Linaria canadensis</i>			
Italian ryegrass	<i>Lolium multiflorum</i>			
Silver-leaf lotus	<i>Lotus argophyllus</i> var. <i>argophyllus</i>			
Grab lotus	<i>Lotus hamatus</i>			
Spanish clover	<i>Lotus purshianus</i> var. <i>purshianus</i>			
Lotus	<i>Lotus salsuginosus</i> var. ?			
Miniature lupine	<i>Lupinus bicolor</i>			
Arroyo lupine	<i>Lupinus succulentus</i>			
Grass poly	<i>Lythrum hyssopifolia</i>			
Laurel sumac	<i>Malosma laurina</i>			
Alkali-mallow	<i>Malvella leprosa</i>			
Man-root	<i>Marah macrocarpus</i> var. <i>macrocarpus</i>			
Horehound	<i>Marrubium vulgare</i>			
Hairy clover fern	<i>Marsilea vestita</i> ssp. <i>vestita</i>			
Pineapple weed	<i>Matricaria matricarioides</i>			
California burclover	<i>Medicago polymorpha</i>			
White sweetclover	<i>Melilotus albus</i>			
Sourclover	<i>Melilotus indicus</i>			
Yellow sweetclover	<i>Melilotus officinalis</i>			
Small-flower microseris	<i>Microseris douglasii</i> ssp. <i>platycarpha</i>			D
Seep monkeyflower	<i>Mimulus guttatus</i>			
Common muilla	<i>Muilla maritima</i>			
Purple needlegrass	<i>Nassella pulchra</i>			
Great marsh evening-primrose	<i>Oenothera elata</i> ssp. <i>hirsutissima</i>			
Prickly-pear	<i>Opuntia</i> sp.			
Osmadenia	<i>Osmadenia tenella</i>			
Kikuyu grass	<i>Pennisetum clandestinum</i>			
Phacelia	<i>Phacelia ramosissima</i> var. ?			
Branching phacelia	<i>Phacelia ramosissima</i> var. <i>latifolia</i>			
Lemmon's canary grass	<i>Phalaris lemmonii</i>			
Paradox canary grass	<i>Phalaris paradoxa</i>			
Bristly ox-tongue	<i>Picris echioides</i>			
American pillwort	<i>Pilularia americana</i>			
Smilo grass	<i>Piptatherum miliaceum</i>			
Adobe popcornflower	<i>Plagiobothrys acanthocarpus</i>			
Rusty popcornflower	<i>Plagiobothrys nothofulvus</i>			
Coast popcornflower	<i>Plagiobothrys undulatus</i>			
Prairie plantain	<i>Plantago elongata</i>			
Dot-seed plantain	<i>Plantago erecta</i>			
Dwarf plantain	<i>Plantago virginica</i>			

Common Name	Scientific Name	Federal	State	County of San Diego
Cream cups	<i>Platystemon californicus</i>			
Annual bluegrass	<i>Poa annua</i>			
Kentucky bluegrass	<i>Poa pratensis</i> ssp. <i>pratensis</i>			
One-sided bluegrass	<i>Poa secunda</i> ssp. <i>secunda</i>			
Common knotweed	<i>Polygonum arenastrum</i>			
Knotweed/ smartweed	<i>Polygonum</i> sp.			
Annual beard grass	<i>Polypogon monspeliensis</i>			
Fremont cottonwood	<i>Populus fremontii</i> ssp. <i>fremontii</i>			
Common purslane	<i>Portulaca oleraceae</i>			
Dwarf woolly-heads	<i>Psilocarphus brevissimus</i> var. <i>brevissimus</i>			
Coast live oak	<i>Quercus agrifolia</i> var. <i>agrifolia</i>			
California buttercup	<i>Ranunculus californicus</i>			
Wild radish	<i>Raphanus sativus</i>			
Skunkbrush	<i>Rhus trilobata</i>			
Water cress	<i>Rorippa nasturtium-aquaticum</i>			
Curly dock	<i>Rumex crispus</i>			
Willow dock	<i>Rumex salicifolius</i> var. ?			
Narrow-leaf willow	<i>Salix exigua</i>			
Goodding's black willow	<i>Salix gooddingii</i>			
Red willow	<i>Salix laevigata</i>			
Arroyo willow	<i>Salix lasiolepis</i>			
Russian thistle	<i>Salsola tragus</i>			
Blue elderberry	<i>Sambucus mexicana</i>			
Sanicle	<i>Sanicula</i> sp.			
Olney's bulrush	<i>Schoenoplectus americanus</i>			
California bulrush	<i>Schoenoplectus californicus</i>			
Common threesquare	<i>Schoenoplectus pungens</i>			
Common threesquare	<i>Schoenoplectus pungens</i>			
California figwort	<i>Scrophularia californica</i> ssp. <i>floribunda</i>			
Bigelow's spike-moss	<i>Selaginella bigelovii</i>			
Checker-bloom	<i>Sidalcea malviflora</i> ssp. <i>sparsifolia</i>			
Common catchfly	<i>Silene gallica</i>			
Milk thistle	<i>Silybum marianum</i>			
Charlock	<i>Sinapis arvensis</i>			
Blue-eyed-grass	<i>Sisyrinchium bellum</i>			
Nightshade	<i>Solanum</i> sp.			
Sow thistle	<i>Sonchus</i> sp.			
Buccone's sand-spurry	<i>Spergularia bocconi</i>			
Common chickweed	<i>Stellaria media</i>			
Salt-cedar	<i>Tamarix</i> sp.			
Puncture vine	<i>Tribulus terrestris</i>			
Vinegar weed	<i>Trichostema lanceolatum</i>			
Tree clover	<i>Trifolium ciliolatum</i>			

Common Name	Scientific Name	Federal	State	County of San Diego
Clover	<i>Trifolium depauperatum</i> var. <i>truncatum</i>			
Strawberry clover	<i>Trifolium fragiferum</i>			
Rose clover	<i>Trifolium hirtum</i>			
Alsike clover	<i>Trifolium hybridum?</i>			
Maiden clover	<i>Trifolium microcephalum</i>			
White tip clover	<i>Trifolium variegatum</i>			
Southern cattail	<i>Typha domingensis</i>			
Hoary nettle	<i>Urtica dioica</i> ssp. <i>holosericea</i>			
Moth mullein	<i>Verbascum blattaria</i>			
Water speedwell	<i>Veronica anagallis-aquatica</i>			
Mexican purslane speedwell	<i>Veronica peregrine</i> ssp. <i>xalapensis</i>			
Purple vetch	<i>Vicia benghalensis</i>			
Narrow-leaf vetch	<i>Vicia sativa</i> ssp. <i>nigra</i>			
Hairy vetch	<i>Vicia villosa</i> ssp. ?			
Johnny-jump-up	<i>Viola pedunculata</i>			
Six-weeks fescue	<i>Vulpia bromoides</i>			
Hairy rat-tail fescue	<i>Vulpia myuros</i> var. <i>hirsuta</i>			
Cocklebur	<i>Xanthium strumarium</i>			
Death camas	<i>Zigadenus venenosus</i> var. <i>venenosus</i>			

FT = Federally Threatened  
SE = California Endangered

## **ATTACHMENT B**

### **ANIMAL SPECIES LIST**

## ATTACHMENT B ANIMAL SPECIES LIST

Common Name	Scientific Name	Federal	State	County of San Diego
<b><i>Invertebrates</i></b>				
Western tiger swallowtail	<i>Papilio rutulus rutulus</i>			
Lorquin's admiral	<i>Basilarchia lorquini</i>			
Buckeye	<i>Junonia coenia</i>			
San Diego fairy shrimp	<i>Branchinecta sandiegonensis</i>	FE		Group 1
Crayfish	<i>Procambarus clarkii</i>			
<b><i>Fishes</i></b>				
Mosquitofish	<i>Gambusia affinis</i>			
Largemouth bass	<i>Micropterus salmoides</i>			
Green sunfish	<i>Lepomis cyanellus</i>			
<b><i>Amphibians</i></b>				
Pacific tree frog	<i>Pseudacris regilla</i>			
Western spadefoot	<i>Spea hammondi</i>		CSC	Group 2
Arroyo toad	<i>Bufo californicus</i>	FE		Group 1
Western toad	<i>Bufo boreas</i>			
Bullfrog	<i>Rana catesbeiana</i>			
<b><i>Reptiles</i></b>				
Two-striped garter snake	<i>Thamnophis hammondi</i>		CSC	Group 1
Gopher snake	<i>Pituophis catenifer annecten</i>			
Coachwhip	<i>Masticophis flagellum</i>			
California kingsnake	<i>Lampropeltis getula</i>			
<b><i>Birds</i></b>				
Great egret	<i>Casmerodius albus</i>			
Snowy egret	<i>Egretta thula</i>			
Canada geese	<i>Branta canadensis</i>			
Mallard	<i>Anas platyrhynchos</i>			
Cinnamon teal	<i>Anas cyanoptera</i>			
Turkey vulture	<i>Cathartes aura</i>			
White-tailed kite	<i>Elanus leucurus</i>		FP	Group 1
Cooper's hawk	<i>Accipiter cooperii</i>		CSC*	Group 1
Red-shouldered hawk	<i>Buteo lineatus</i>			Group 1
Red-tailed hawk	<i>Buteo jamaicensis</i>			
Zone-tailed hawk	<i>Buteo abonotatus</i>			
Ferruginous hawk	<i>Buteo regalis</i>		CSC**	Group 1**
Northern harrier	<i>Circus cyaneus</i>		CSC*	Group 1
American kestrel	<i>Falco sparverius</i>			
Merlin	<i>Falco columbarius</i>		CSC**	Group 2**
Prairie falcon	<i>Falco mexicanus</i>		CSC*	Group 1
Golden eagle	<i>Aquila chrysaetos</i>		CSC, FP	Group 1
Bald eagle	<i>Haliaeetus leucocephalus</i>		SE, FP	Group 1**
Barn owl	<i>Tyto alba</i>			Group 2
Great-horned owl	<i>Bubo virginianus</i>			

Common Name	Scientific Name	Federal	State	County of San Diego
Burrowing owl	<i>Athene cunicularia</i>		CSC*	Group 1
Virginia rail	<i>Rallus limicola</i>			
Sora	<i>Porzana carolina</i>			
Killdeer	<i>Charadrius vociferus</i>			
Mourning dove	<i>Zenaida macroura</i>			
Anna's hummingbird	<i>Calypte anna</i>			
Acorn woodpecker	<i>Melanerpes formicivorus</i>			
Nuttall's woodpecker	<i>Picoides nuttallii</i>			
Northern flicker	<i>Colaptes cafer</i>			
Western wood-pewee	<i>Contopus sordidulus</i>			
Willow flycatcher	<i>Empidonax traillii estimus***</i>	FE*	SE*	Group 1
Black phoebe	<i>Sayornis nigricans</i>			
Say's phoebe	<i>Sayornis saya</i>			
Ash-throated flycatcher	<i>Myiarchus cinerascens</i>			
Cassin's kingbird	<i>Tyrannus vociferans</i>			
Western kingbird	<i>Tyrannus verticalis</i>			
Loggerhead shrike	<i>Lanius ludovicianus</i>		CSC*	Group 1
Warbling vireo	<i>Vireo gilvus</i>			
Western scrub-jay	<i>Aphelocoma californica</i>			
American crow	<i>Corvus brachyrhynchos</i>			
Common raven	<i>Corvus corax</i>			
N. rough-winged swallow	<i>Stelgidopteryx serripennis</i>			
Oak titmouse	<i>Baeolophus inornatus</i>			
Bushtit	<i>Psaltiriparus minimus</i>			
Bewick's wren	<i>Thryomanes bewickii</i>			
House wren	<i>Troglodytes aedon</i>			
Coastal California gnatcatcher	<i>Poliophtila californica californica</i>	FT		Group 1
Western bluebird	<i>Sialia mexicana</i>			Group 2
Northern mockingbird	<i>Mimus polyglottos</i>			
European starling	<i>Sturnus vulgaris</i>			
Yellow warbler	<i>Dendroica petechia</i>		CSC	Group 2
Common yellowthroat	<i>Geothlypis trichas</i>			
Spotted towhee	<i>Pipilo maculatus</i>			
California towhee	<i>Pipilo crissalis</i>			
Lark sparrow	<i>Chondestes grammacus</i>			
Grasshopper sparrow	<i>Ammodramus savannarum</i>			Group 1
Song sparrow	<i>Melospiza melodia</i>			
Black-headed grosbeak	<i>Pheucticus melanocephalus</i>			
Blue grosbeak	<i>Guiraca caerulea</i>			

Common Name	Scientific Name	Federal	State	County of San Diego
Red-winged blackbird	<i>Agelaius phoenicius</i>			
Western meadowlark	<i>Sturnella neglecta</i>			
Brown-headed cowbird	<i>Molothrus ater</i>			
Hooded oriole	<i>Icterus cucullatus</i>			
Bullock's oriole	<i>Icterus galbula</i>			
House finch	<i>Carpodacus mexicanus</i>			
Lesser goldfinch	<i>Carduelis psaltria</i>			
American goldfinch	<i>Carduelis tristis</i>			
House sparrow	<i>Passer domesticus</i>			
<b>Mammals</b>				
Botta's pocket gopher	<i>Thomomys bottae</i>			
Dulzura kangaroo rat	<i>Dipodomys simulans</i>			
Stephens' kangaroo rat	<i>Dipodomys stephensi</i>	<b>FE</b>	<b>ST</b>	<b>Group 1</b>
Audubon's cottontail	<i>Sylvilagus audubonii</i>			
California ground squirrel	<i>Spermophilus beecheyi</i>			
Long-tailed weasel	<i>Mustela frenata</i>			
Gray fox	<i>Urocyon cinereoargenteus</i>			
Coyote	<i>Canis latrans</i>			
Mule deer	<i>Odocoileus hemionus</i>			<b>Group 2</b>
Bobcat	<i>Felis rufus</i>			
Mountain lion	<i>Felis concolor</i>			<b>Group 2</b>

FE = Federally Endangered

FT = Federally Threatened

SE = California Endangered

ST = California Threatened

CSC = California Species of Special Concern

FP = California Fully Protected

\* = Nesting locations

\*\* = Wintering locations

\*\*\* = A single willow flycatcher was detected on a single date and was not confirmed to subspecies.

**ATTACHMENT C  
PHOTOGRAPHS OF  
SELECTED SENSITIVE SPECIES**

## **ATTACHMENT C**

### **PHOTOGRAPHS OF SELECTED SENSITIVE SPECIES**



Figure C1. Stephens' kangaroo rat (SKR) burrow (top) and individual (bottom). Photos by Michael White and Wayne Spencer.



Figure C2. Ferruginous hawk (top) and golden eagle (bottom). Photos by WRI and George Hartwell.



Figure C3. Alkali playa bordered by southern tarplant (top) and southern tarplant (bottom). Photos by Michael White



Figure C4. Parish's brittlescale (top) and Coulter's saltbush (bottom). Photos by Michael White.



Figure C5. Arroyo toad habitat in Reach A of Santa Maria Creek (top) and arroyo toad (bottom).  
Photos by Michael White



Figure C6. (Top) Small-leaved morning glory (left) and round-leaved filaree (right) and (bottom) small-leaved morning glory. Photos by Gena Calcarone.

# **ATTACHMENT D**

# **PHOTOMONITORING RESULTS**

## ATTACHMENT D

### PHOTOMONITORING RESULTS



Figure D1. Grassland plot 14 looking south in 2005 (top) and 2006 (bottom).



Figure D2. Vernal pool e44 looking north in 2005 (top) and 2006 (bottom).



Figure D3. Riparian transect 3 looking south in 2005 (top) and 2006 (bottom).



Figure D4. Riparian transect 5 looking south in 2005 (top) and 2006 (bottom).



Figure D5. Riparian transect 10 looking south in 2005 (top) and 2006 (bottom).

# **ATTACHMENT E**

## **GRASSLAND SPECIES RICHNESS**

## ATTACHMENT E GRASSLAND SPECIES RICHNESS

SPECIES	Grassland Plot #s																								Species Total
	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	
<i>Achyrachaena mollis</i>				X																					1
<i>Ambrosia psilostachya</i>	X		X	X		X	X	X	X	X		X	X	X	X		X	X	X	X		X			17
<i>Amsinckia</i> sp.		X							X																2
<i>Anagallis arvensis</i>				X													X								2
<i>Anthemis cotula</i>			X		X	X	X	X	X					X			X	X				X			10
<i>Asclepias californica</i>							X	X	X	X					X						X	X			7
<i>Atriplex parishii</i> var. <i>parishii</i>																	X								1
<i>Avena barbata</i>	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	24
<i>Avena fatua</i>																								X	1
<i>Brassica nigra</i>			X																					X	2
<i>Brickellia californica</i>									X																1
<i>Brodiaea terrestris</i> ssp. <i>kernensis</i>														X						X					2
<i>Bromus diandrus</i>	X		X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	23
<i>Bromus hordeaceus</i>	X	X	X	X	X	X	X	X	X	X	X	X	X	X		X	X	X	X	X	X	X	X	X	23
<i>Bromus madritensis</i> ssp. <i>rubens</i>			X	X	X				X	X	X			X	X	X					X				10

SPECIES	Grassland Plot #s																								Species Total
	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	
<i>Calochortus splendens</i>		X		X										X		X				X					5
<i>Calystegia macrostegia</i>		X												X											2
<i>Camissonia claviformis</i>					X																				1
<i>Camissonia strigulosa</i>					X				X	X	X														4
<i>Castilleja</i> sp.																			X						1
<i>Centaurea melitensis</i>		X												X		X								X	4
<i>Centaureum venustum</i>														X			X								2
<i>Centromadia parryi</i> ssp. <i>australis</i>														X			X	X				X			4
<i>Chlorogalum parviflorum</i>		X																						X	2
<i>Clarkia purpurea</i>							X			X	X														3
<i>Convolvulus simulans</i>			X																					X	2
<i>Conyza</i> sp.					X				X	X				X			X	X							6
<i>Corethrogyne filaginifolia</i> var. <i>filaginifolia</i>						X	X			X						X				X	X				6
<i>Cotula coronopifolia</i>																			X						1
<i>Crassula connata</i>														X					X						2
<i>Croton setigerus</i>	X	X	X	X	X	X		X	X	X	X	X		X	X	X	X	X	X	X	X	X			20
<i>Cryptantha</i> sp.				X							X					X				X					4

SPECIES	Grassland Plot #s																								Species Total
	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	
<i>Cucurbita palmata</i>					X																				1
<i>Cynara cardunculus</i>			X											X										X	3
<i>Cynodon dactylon</i>						X											X	X				X			4
<i>Daucus pusillus</i>		X																							1
<i>Deinandra fasciculata</i>		X	X	X	X	X			X				X	X		X	X	X	X	X		X	X	X	16
<i>Distichlis spicata</i>	X	X	X	X	X	X	X	X	X	X	X	X	X		X	X	X	X	X	X	X	X			21
<i>Eleocharis macrostachya</i>			X																						1
<i>Eriogonum elongatum</i> var. <i>elongatum</i>		X																							1
<i>Eriogonum fasciculatum</i> var. <i>fasciculatum</i>		X									X			X											3
<i>Eriogonum fasciculatum</i> var. <i>polifolium</i>									X		X											X			3
<i>Erodium botrys</i>	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X		X	23
<i>Erodium cicutarium</i>			X																						1
<i>Euphorbia spathulata</i>																								X	1
<i>Filago gallica</i>			X											X											2
<i>Gastroidium ventricosum</i>						X	X	X	X	X	X			X					X		X				9
<i>Gilia</i> sp.										X															1

SPECIES	Grassland Plot #s																								Species Total	
	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24		
<i>Gnaphalium californicum</i>									X	X	X															3
<i>Grindelia camporum</i> var. <i>bracteosa</i>	X					X										X			X	X		X	X	X		8
<i>Hazardia squarrosa</i> var. <i>grindelioides</i>				X		X	X		X																	4
<i>Hedypnois cretica</i>			X																							1
<i>Heliotropium curassavicum</i>				X																		X				2
<i>Hirschfeldia incana</i>		X	X		X		X	X	X	X	X	X	X	X	X	X	X	X	X		X	X	X	X		20
<i>Holocarpha virgata</i> ssp. <i>elongata</i>	X		X	X	X							X	X						X							7
<i>Hordeum marinum</i> ssp. <i>gussoneanum</i>	X	X	X	X	X	X	X	X	X		X		X	X	X	X	X	X	X	X		X	X	X		21
<i>Hordeum murinum</i> ssp. <i>leporinum</i>					X																	X				2
<i>Hypochaeris glabra</i>			X	X	X	X					X			X			X			X						8
<i>Isocoma menziesii</i> var. <i>menziesii</i>		X	X	X		X									X	X			X	X	X	X				10
<i>Juncus bufonius</i> var. <i>bufonius</i>			X	X		X	X	X		X				X			X	X	X							10
<i>Lactuca serriola</i>			X	X					X									X								4

SPECIES	Grassland Plot #s																								Species Total
	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	
<i>Lamarckia aurea</i>		X							X	3	X														6
<i>Leymus triticoides</i>					X												X	X							3
<i>Linaria canadensis</i>								X		X	X									X					4
<i>Lolium multiflorum</i>	X	X	X	X	X	X	X	X	X	X	X		X	X		X	X	X	X	X	X	X	X	X	22
<i>Lotus argophyllus</i> var. <i>argophyllus</i>									X	X	X														3
<i>Lotus hamatus</i>		X	X								X														3
<i>Lupinus bicolor</i>			X		X	X	X	X	X	X							X	X		X					10
<i>Lupinus succulentus</i>																X									1
<i>Lythrum hyssopifolia</i>			X			X		X					X				X	X	X	X					8
<i>Malosma laurina</i>									X																1
<i>Medicago polymorpha</i>			X	X												X	X	X	X						6
<i>Melilotus indicus</i>																	X		X						2
<i>Microseris douglasii</i> ssp. <i>platycarpa</i>												X											X		2
<i>Muilla maritima</i>	X											X	X						X	X					5
<i>Nassella pulchra</i>		X	X	X							X	X	X	X					X					X	9
<i>Osmadenia tenella</i>		X																							1

SPECIES	Grassland Plot #s																								Species Total
	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	
<i>Phacelia ramosissima</i> var. <i>latifolia</i>									X		X														2
<i>Phalaris paradoxa</i>			X																					X	2
<i>Plantago erecta</i>				X									X	X		X							X	X	6
<i>Plantago</i> sp.									X																1
<i>Plantago virginica</i>			X																					X	2
<i>Polygonum arenastrum</i>																			X						1
<i>Polypogon monspeliensis</i>						X											X		X			X			4
<i>Psilocarphus brevissimus</i> var. <i>brevissimus</i>																				X					1
<i>Quercus agrifolia</i> var. <i>agrifolia</i>																	X								1
<i>Raphanus sativus</i>														X											1
<i>Rumex crispus</i>					X		X							X			X	X	X	X		X			8
<i>Rumex salicifolius</i>	X		X	X			X									X			X						6
<i>Sambucus mexicana</i>									X		X														2
<i>Scrophularia californica</i> ssp. <i>floribunda</i>									X	X	X														3
<i>Silene gallica</i>	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X			22
<i>Solanum</i> sp.		X																							1
<i>Sonchus</i> sp.														X											1

SPECIES	Grassland Plot #s																								Species Total
	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	
<i>Spergularia bocconi</i>																	X		X						2
<i>Stellaria media</i>						X																			1
<i>Trichostema lanceolatum</i>		X												X					X	X					4
<i>Trifolium ciliolatum</i>								X																	1
<i>Trifolium depauperatum</i> var. <i>truncatum</i>			X	X																					2
<i>Trifolium fragiferum</i>																		X							1
<i>Trifolium hirtum</i>		X	X							X															3
<i>Trifolium microcephalum</i>			X								X														2
<i>Trifolium</i> sp.										X							X	X							3
<i>Verbascum blattaria</i>																	X								1
<i>Veronica peregrina</i> ssp. <i>xalapensis</i>																			X						1
<i>Vicia sativa</i> ssp. <i>nigra</i>																	X								1
<i>Vicia villosa</i>														X											1
<i>Vulpia myuros</i> var. <i>hirsuta</i>	X	X	X	X		X	X	X		X	X	X	X	X	X	X	X	X	X	X	X	X			20
<i>Xanthium strumarium</i>																						X			1
<b>Plot Totals</b>	15	27	38	29	23	25	21	20	32	30	29	14	16	36	13	23	34	26	34	25	15	25	9	21	580

# **ATTACHMENT F**

## **VERNAL POOL SPECIES RICHNESS**

# ATTACHMENT F VERNAL POOL SPECIES RICHNESS

Species	Airport Vernal Pools								Cagney Vernal Pools				Cumming Swale		Cagney Swale				Alkali Playa RAAP 100	Total Frequency of Occurrence
	e44	e45	e46	e52	e53	e54	e77	r24	e56	e58	e59	e62	ev1	ev1	VS1	VS1	VS3	VS4		
<i>Amaranthus</i> sp.					X															1
<i>Ambrosia psilostachya</i>	X	X		X	X	X	X	X			X	X	X	X	X	X	X	X	X	13
<i>Amsinckia</i> sp.					X															1
<i>Anagallis arvensis</i>															X	X			X	3
<i>Anemopsis californica</i>														X						1
<i>Anthemis cotula</i>									X	X	X	X	X	X	X	X	X	X	X	11
<i>Avena barbata</i>	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	19
<i>Brassica nigra</i>						X							X							2
<i>Brodiaea terrestris</i> ssp. <i>kernensis</i>	X				X	X		X	X			X		X			X			8
<i>Bromus diandrus</i>		X	X	X	X	X	X	X	X	X		X	X	X	X	X	X	X	X	16
<i>Bromus hordeaceus</i>	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	19
<i>Bromus madritensis</i> ssp. <i>rubens</i>			X					X							X				X	4
<i>Calandrinia ciliata</i>										X										1
<i>Callitriche</i> sp.								X					X		X	X	X	X		5
<i>Calochortus splendens</i>					X															1
<i>Castilleja densiflora</i> ssp. <i>densiflora</i>	X	X	X			X		X				X								5
<i>Centromadia parryi</i> ssp. <i>australis</i>									X	X	X			X	X	X	X	X	X	9

Species	Airport Vernal Pools								Cagney Vernal Pools					Cumming Swale		Cagney Swale				Alkali Playa RAAP 100	Total Frequency of Occurrence
	e44	e45	e46	e52	e53	e54	e77	r24	e56	e58	e59	e62	ev1	ev1	VS1	VS1	VS3	VS4			
<i>Centunculus minimus</i>											X		X	X			X	X			5
<i>Cerastium glomeratum</i>										X											1
<i>Cotula coronopifolia</i>									X	X	X	X	X	X	X	X	X	X			9
<i>Crassula aquatica</i>	X		X	X	X			X	X	X	X	X	X	X	X	X	X	X			15
<i>Croton setigerus</i>		X													X			X			3
<i>Crypsis schoenoides</i>																					1
<i>Cryptantha</i> sp.		X				X		X						X							3
<i>Cynodon dactylon</i>									X	X		X	X	X	X	X	X	X		X	10
<i>Deinandra fasciculata</i>	X	X	X	X	X	X	X	X	X	X	X	X	X			X	X	X	X		16
<i>Deschampsia danthonioides</i>			X																		1
<i>Dichelostemma capitatum</i> ssp. <i>capitatum</i>																					
<i>Distichlis spicata</i>	X	X	X	X		X	X		X	X	X	X	X	X	X	X	X	X	X	X	17
<i>Downingia cuspidata</i>			X																		1
<i>Eleocharis macrostachya</i>	X	X	X	X	X	X		X	X	X	X	X	X	X	X	X	X	X	X	X	18
<i>Elyrigia</i> sp.																					1
<i>Epilobium pygmaeum</i>			X																		1
<i>Eremocarpus setigerus</i>		X														X	X	X			4
<i>Erodium botrys</i>	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	19
<i>Grindelia camporum</i> var. <i>bracteosum</i>								X													
<i>Heliotropium curassavicum</i>						X	X	X								X		X			4
										</											

Species	Airport Vernal Pools								Cagney Vernal Pools				Cumming Swale				Cagney Swale				Alkali Playa RAAP 100	Total Frequency of Occurrence
	e44	e45	e46	e52	e53	e54	e77	r24	e56	e58	e59	e62	ev1	ev1	VS1	VS1	VS3	VS4				
<i>Hirschfeldia incana</i>									X		X		X	X	X						X	6
<i>Hordeum marinum</i> ssp. <i>gussonianum</i>	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	19
<i>Hordeum murinum</i> ssp. <i>leporinum</i>																X		X				2
<i>Hypochaeris glabra</i>		X	X	X	X	X			X			X						X				8
<i>Isocoma menziesii</i> var. <i>menziesii</i>				X																		1
<i>Isoetes howellii</i>																	X					1
<i>Juncus bufonius</i> var. <i>bufonius</i>	X	X	X	X	X	X	X	X	X	X	X	X			X	X	X	X	X	X	X	17
<i>Juncus mexicanus</i>																X						1
<i>Juncus</i> sp.		X																				1
<i>Juncus xiphioides</i>		X		X		X	X			X			X			X		X		X	X	9
<i>Lepidium nitidum</i> var. <i>nitidum</i>	X					X				X												3
<i>Lepidium oblongum</i> var. <i>insulare</i>																				X		1
<i>Lessingia filaginifolia</i> var. <i>filaginifolia</i>		X	X	X		X																4
<i>Leymus triticoides</i>												X		X				X				3
<i>Lilaea scilloides</i>									X		X				X		X	X				5
<i>Limosella acutis</i>											X				X	X	X	X				4
<i>Lolium multiflorum</i>	X		X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	18
<i>Lotus hamatus</i>			X		X																	2
<i>Lotus purshianus</i> var. <i>purshianus</i>				X				X										X				3
<i>Lupinus bicolor</i>		X	X	X	X	X	X	X		X		X			X	X	X	X	X			12

Species	Airport Vernal Pools								Cagney Vernal Pools				Cumming Swale		Cagney Swale				Alkali Playa	Total Frequency of Occurrence	
	e44	e45	e46	e52	e53	e54	e77	r24	e56	e58	e59	e62	ev1	ev1	VS1	VS1	VS3	VS4	RAAP 100		
<i>Lythrum hyssopifolium</i>	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	19
<i>Malvella leprosa</i>									X					X					X	3	
<i>Marsilea vestita</i> ssp. <i>vestita</i>											X		X	X			X			5	
<i>Medicago polymorpha</i>	X					X		X		X	X	X	X	X	X	X	X	X	X	12	
<i>Melilotus indica</i>															X					1	
<i>Melilotus</i> sp.																	X	X	X	3	
<i>Microseris douglasii</i> ssp. <i>platycarpa</i>	X							X												2	
<i>Muilla maritima</i>		X	X	X	X	X	X													6	
<i>Pilularia americana</i>											X									1	
<i>Plagiobothrys acanthocarpus</i>	X																			1	
<i>Plagiobothrys</i> sp.							X													1	
<i>Plagiobothrys undulatus</i>	X		X		X			X	X	X	X	X				X	X	X		10	
<i>Plantago elongata</i>						X	X					X								3	
<i>Plantago erecta</i>	X					X	X	X												4	
<i>Poa annua</i>										X							X			2	
<i>Polygonum arenastrum</i>								X	X		X					X	X	X	X	6	
<i>Polypogon monspeliensis</i>								X	X		X		X	X	X	X	X	X	X	9	
<i>Psilocarphus brevissimus</i> var. <i>brevissimus</i>	X	X	X	X	X	X		X	X	X	X	X	X	X	X	X	X	X		17	
<i>Rumex crispus</i>			X		X			X	X	X	X	X	X	X	X	X	X	X	X	14	
<i>Rumex salicifolius</i>									X	X								X	X	4	
<i>Salsola tragus</i>								X												1	
<i>Sanicula</i> sp.		X																		1	
<i>Silene gallica</i>			X		X	X			X	X			X	X	X	X	X	X	X	11	

Species	Airport Vernal Pools								Cagney Vernal Pools						Cumming Swale		Cagney Swale					Alkali Playa	Total Frequency of Occurrence
	e44	e45	e46	e52	e53	e54	e77	r24	e56	e58	e59	e62	ev1	ev1	VS1	VS1	VS3	VS4	RAAP 100				
<i>Sisyrinchium bellum</i>		X	X			X															3		
<i>Sonchus</i> sp.		X							X												2		
<i>Spergularia bocconii</i>										X		X				X					3		
<i>Trichostema lanceolatum</i>	X	X	X	X	X	X									X						7		
<i>Trifolium ciliolatum</i>					X																1		
<i>Trifolium depauperatum</i> var. <i>truncatum</i>	X	X	X	X	X	X	X		X												8		
<i>Trifolium</i> sp.				X					X	X											3		
<i>Trifolium variegatum</i>									X												1		
<i>Typha</i> sp.														X							1		
<i>Veronica anagallis-aquatica</i>													X			X	X	X	X		4		
<i>Veronica peregrina</i> ssp. <i>xalapensis</i>	X		X			X		X	X	X	X	X				X	X	X			10		
<i>Vicia villosa</i>													X								1		
<i>Vulpia bromoides</i>					X																1		
<i>Vulpia myuros</i> var. <i>hirsuta</i>	X	X		X	X	X	X	X	X	X	X	X			X			X	X		14		
<i>Xanthium strumarium</i>														X	X	X	X	X			5		
Total Species	25	28	30	24	30	33	21	31	34	31	28	29	27	28	30	36	35	42	29		570		

# **ATTACHMENT G**

## **WATER QUALITY RESULTS**

## ATTACHMENT G WATER QUALITY RESULTS

Sample Date	Station	Source Description	Analysis	Analyte name	Value	MDL*	Units
3-Jan-05	SMC1	Upstream sample site Ramona Grasslands	NH3_NO3_NO2_AUT	Ammonia-N	0.099	0.04	MG/L
24-Jan-05	SMC1	Upstream sample site Ramona Grasslands	NH3_NO3_NO2_AUT	Ammonia-N	ND	0.04	MG/L
14-Feb-05	SMC1	Upstream sample site Ramona Grasslands	NH3_NO3_NO2_AUT	Ammonia-N	0.348	0.04	MG/L
28-Feb-05	SMC1	Upstream sample site Ramona Grasslands	NH3_NO3_NO2_AUT	Ammonia-N	0.476	0.04	MG/L
15-Mar-05	SMC1	Upstream sample site Ramona Grasslands	NH3_NO3_NO2_AUT	Ammonia-N	0.049	0.04	MG/L
28-Mar-05	SMC1	Upstream sample site Ramona Grasslands	NH3_NO3_NO2_AUT	Ammonia-N	ND	0.04	MG/L
11-Apr-05	SMC1	Upstream sample site Ramona Grasslands	NH3_NO3_NO2_AUT	Ammonia-N	ND	0.04	MG/L
25-Apr-05	SMC1	Upstream sample site Ramona Grasslands	NH3_NO3_NO2_AUT	Ammonia-N	ND	0.04	MG/L
9-May-05	SMC1	Upstream sample site Ramona Grasslands	NH3_NO3_NO2_AUT	Ammonia-N	ND	0.04	MG/L
23-May-05	SMC1	Upstream sample site Ramona Grasslands	NH3_NO3_NO2_AUT	Ammonia-N	ND	0.04	MG/L
13-Jun-05	SMC1	Upstream sample site Ramona Grasslands	NH3_NO3_NO2_AUT	Ammonia-N	ND	0.04	MG/L
13-Mar-06	SMC1	Upstream sample site Ramona Grasslands	NH3_NO3_NO2_AUT	Ammonia-N	ND	0.031	MG/L
27-Mar-06	SMC1	Upstream sample site Ramona Grasslands	NH3_NO3_NO2_AUT	Ammonia-N	ND	0.031	MG/L
3-Jan-05	SMC1	Upstream sample site Ramona Grasslands	ANIONS_IC	Bromide	0.131	0.1	MG/L
28-Feb-05	SMC1	Upstream sample site Ramona Grasslands	ANIONS_IC	Bromide	0.224	0.1	MG/L
11-Apr-05	SMC1	Upstream sample site Ramona Grasslands	ANIONS_IC	Bromide	0.384	0.1	MG/L
25-Apr-05	SMC1	Upstream sample site Ramona Grasslands	ANIONS_IC	Bromide	0.359	0.1	MG/L

Sample Date	Station	Source Description	Analysis	Analyte name	Value	MDL*	Units
9-May-05	SMC1	Upstream sample site Ramona Grasslands	ANIONS_IC	Bromide	0.399	0.1	MG/L
23-May-05	SMC1	Upstream sample site Ramona Grasslands	ANIONS_IC	Bromide	0.573	0.1	MG/L
13-Jun-05	SMC1	Upstream sample site Ramona Grasslands	ANIONS_IC	Bromide	0.676	0.1	MG/L
13-Mar-06	SMC1	Upstream sample site Ramona Grasslands	ANIONS_IC	Bromide	0.405	0.1	MG/L
27-Mar-06	SMC1	Upstream sample site Ramona Grasslands	ANIONS_IC_LOW	Bromide	0.52	0.025	MG/L
3-Jan-05	SMC1	Upstream sample site Ramona Grasslands	ANIONS_IC	Chloride	57.1	0.5	MG/L
28-Feb-05	SMC1	Upstream sample site Ramona Grasslands	ANIONS_IC	Chloride	90.6	1	MG/L
11-Apr-05	SMC1	Upstream sample site Ramona Grasslands	ANIONS_IC	Chloride	148	1	MG/L
25-Apr-05	SMC1	Upstream sample site Ramona Grasslands	ANIONS_IC	Chloride	152	1	MG/L
9-May-05	SMC1	Upstream sample site Ramona Grasslands	ANIONS_IC	Chloride	150	1	MG/L
23-May-05	SMC1	Upstream sample site Ramona Grasslands	ANIONS_IC	Chloride	190	1	MG/L
13-Jun-05	SMC1	Upstream sample site Ramona Grasslands	ANIONS_IC	Chloride	228	1	MG/L
13-Mar-06	SMC1	Upstream sample site Ramona Grasslands	ANIONS_IC	Chloride	154	0.5	MG/L
27-Mar-06	SMC1	Upstream sample site Ramona Grasslands	ANIONS_IC_LOW	Chloride	200	0.125	MG/L
3-Jan-05	SMC1	Upstream sample site Ramona Grasslands	STREAM_PARAM	Conductivity	479		US/CM
24-Jan-05	SMC1	Upstream sample site Ramona Grasslands	STREAM_PARAM	Conductivity	1610		US/CM
14-Feb-05	SMC1	Upstream sample site Ramona Grasslands	STREAM_PARAM	Conductivity	1230		US/CM
28-Feb-05	SMC1	Upstream sample site Ramona Grasslands	STREAM_PARAM	Conductivity	854		US/CM
15-Mar-05	SMC1	Upstream sample site Ramona Grasslands	STREAM_PARAM	Conductivity	1120		US/CM

Sample Date	Station	Source Description	Analysis	Analyte name	Value	MDL*	Units
28-Mar-05	SMC1	Upstream sample site Ramona Grasslands	STREAM_PARAM	Conductivity	1090		US/CM
11-Apr-05	SMC1	Upstream sample site Ramona Grasslands	STREAM_PARAM	Conductivity	1250		US/CM
25-Apr-05	SMC1	Upstream sample site Ramona Grasslands	STREAM_PARAM	Conductivity	1220		US/CM
9-May-05	SMC1	Upstream sample site Ramona Grasslands	STREAM_PARAM	Conductivity	1230		US/CM
23-May-05	SMC1	Upstream sample site Ramona Grasslands	STREAM_PARAM	Conductivity	1430		US/CM
13-Jun-05	SMC1	Upstream sample site Ramona Grasslands	STREAM_PARAM	Conductivity	1600		US/CM
13-Mar-06	SMC1	Upstream sample site Ramona Grasslands	STREAM_PARAM	Conductivity	1030		US/CM
3-Jan-05	SMC1	Upstream sample site Ramona Grasslands	STREAM_PARAM	Dissolved Oxygen	9.76		MG/L
24-Jan-05	SMC1	Upstream sample site Ramona Grasslands	STREAM_PARAM	Dissolved Oxygen	10.1		MG/L
14-Feb-05	SMC1	Upstream sample site Ramona Grasslands	STREAM_PARAM	Dissolved Oxygen	9.13		MG/L
28-Feb-05	SMC1	Upstream sample site Ramona Grasslands	STREAM_PARAM	Dissolved Oxygen	10.1		MG/L
15-Mar-05	SMC1	Upstream sample site Ramona Grasslands	STREAM_PARAM	Dissolved Oxygen	10.8		MG/L
28-Mar-05	SMC1	Upstream sample site Ramona Grasslands	STREAM_PARAM	Dissolved Oxygen	10.3		MG/L
11-Apr-05	SMC1	Upstream sample site Ramona Grasslands	STREAM_PARAM	Dissolved Oxygen	10.6		MG/L
25-Apr-05	SMC1	Upstream sample site Ramona Grasslands	STREAM_PARAM	Dissolved Oxygen	10.4		MG/L
9-May-05	SMC1	Upstream sample site Ramona Grasslands	STREAM_PARAM	Dissolved Oxygen	9.09		MG/L
23-May-05	SMC1	Upstream sample site Ramona Grasslands	STREAM_PARAM	Dissolved Oxygen	6.99		MG/L
13-Jun-05	SMC1	Upstream sample site Ramona Grasslands	STREAM_PARAM	Dissolved Oxygen	12.3		MG/L
13-Mar-06	SMC1	Upstream sample site Ramona Grasslands	STREAM_PARAM	Dissolved Oxygen	10.6		MG/L

Sample Date	Station	Source Description	Analysis	Analyte name	Value	MDL*	Units
3-Jan-05	SMC1	Upstream sample site Ramona Grasslands	QT_CST	Enterococcus	> 2400		/100 ML
24-Jan-05	SMC1	Upstream sample site Ramona Grasslands	QT_CST	Enterococcus	28		/100 ML
14-Feb-05	SMC1	Upstream sample site Ramona Grasslands	QT_CST	Enterococcus	310		/100 ML
28-Feb-05	SMC1	Upstream sample site Ramona Grasslands	QT_CST	Enterococcus	110		/100 ML
15-Mar-05	SMC1	Upstream sample site Ramona Grasslands	QT_CST	Enterococcus	170		/100 ML
28-Mar-05	SMC1	Upstream sample site Ramona Grasslands	QT_CST	Enterococcus	47		/100 ML
11-Apr-05	SMC1	Upstream sample site Ramona Grasslands	QT_CST	Enterococcus	80		/100 ML
25-Apr-05	SMC1	Upstream sample site Ramona Grasslands	QT_CST	Enterococcus	100		/100 ML
9-May-05	SMC1	Upstream sample site Ramona Grasslands	QT_CST	Enterococcus	84		/100 ML
23-May-05	SMC1	Upstream sample site Ramona Grasslands	QT_CST	Enterococcus	1100		/100 ML
13-Jun-05	SMC1	Upstream sample site Ramona Grasslands	QT_CST	Enterococcus	190		/100 ML
13-Mar-06	SMC1	Upstream sample site Ramona Grasslands	QT_CST	Enterococcus	330		/100 ML
27-Mar-06	SMC1	Upstream sample site Ramona Grasslands	QT_CST	Enterococcus	110		/100 ML
3-Jan-05	SMC1	Upstream sample site Ramona Grasslands	QT_CST	Escherichia Coli	3400		/100 ML
24-Jan-05	SMC1	Upstream sample site Ramona Grasslands	QT_CST	Escherichia Coli	410		/100 ML
14-Feb-05	SMC1	Upstream sample site Ramona Grasslands	QT_CST	Escherichia Coli	520		/100 ML
28-Feb-05	SMC1	Upstream sample site Ramona Grasslands	QT_CST	Escherichia Coli	520		/100 ML
15-Mar-05	SMC1	Upstream sample site Ramona Grasslands	QT_CST	Escherichia Coli	200		/100 ML
28-Mar-05	SMC1	Upstream sample site Ramona Grasslands	QT_CST	Escherichia Coli	100		/100 ML

Sample Date	Station	Source Description	Analysis	Analyte name	Value	MDL*	Units
11-Apr-05	SMC1	Upstream sample site Ramona Grasslands	QT_CST	Escherichia Coli	520		/100 ML
25-Apr-05	SMC1	Upstream sample site Ramona Grasslands	QT_CST	Escherichia Coli	200		/100 ML
9-May-05	SMC1	Upstream sample site Ramona Grasslands	QT_CST	Escherichia Coli	520		/100 ML
23-May-05	SMC1	Upstream sample site Ramona Grasslands	QT_CST	Escherichia Coli	310		/100 ML
13-Jun-05	SMC1	Upstream sample site Ramona Grasslands	QT_CST	Escherichia Coli	310		/100 ML
13-Mar-06	SMC1	Upstream sample site Ramona Grasslands	QT_CST	Escherichia Coli	410		/100 ML
27-Mar-06	SMC1	Upstream sample site Ramona Grasslands	QT_CST	Escherichia Coli	100		/100 ML
3-Jan-05	SMC1	Upstream sample site Ramona Grasslands	NH3_NO3_NO2_AUT	Nitrate	6.76	0.3	MG/L
24-Jan-05	SMC1	Upstream sample site Ramona Grasslands	NH3_NO3_NO2_AUT	Nitrate	19.4	0.6	MG/L
14-Feb-05	SMC1	Upstream sample site Ramona Grasslands	NH3_NO3_NO2_AUT	Nitrate	19.5	0.6	MG/L
28-Feb-05	SMC1	Upstream sample site Ramona Grasslands	NH3_NO3_NO2_AUT	Nitrate	31.6	0.6	MG/L
15-Mar-05	SMC1	Upstream sample site Ramona Grasslands	NH3_NO3_NO2_AUT	Nitrate	42.9	1.2	MG/L
28-Mar-05	SMC1	Upstream sample site Ramona Grasslands	NH3_NO3_NO2_AUT	Nitrate	37.3	0.6	MG/L
11-Apr-05	SMC1	Upstream sample site Ramona Grasslands	ANIONS_IC	Nitrate	46.7	0.6	MG/L
25-Apr-05	SMC1	Upstream sample site Ramona Grasslands	NH3_NO3_NO2_AUT	Nitrate	> 25	0.3	MG/L
9-May-05	SMC1	Upstream sample site Ramona Grasslands	ANIONS_IC	Nitrate	38.2	0.2	MG/L
9-May-05	SMC1	Upstream sample site Ramona Grasslands	NH3_NO3_NO2_AUT	Nitrate	35.9	0.6	MG/L
23-May-05	SMC1	Upstream sample site Ramona Grasslands	NH3_NO3_NO2_AUT	Nitrate	16.6	0.6	MG/L
13-Jun-05	SMC1	Upstream sample site Ramona Grasslands	NH3_NO3_NO2_AUT	Nitrate	0.902	0.06	MG/L

Sample Date	Station	Source Description	Analysis	Analyte name	Value	MDL*	Units
13-Mar-06	SMC1	Upstream sample site Ramona Grasslands	ANIONS_IC	Nitrate	1.91	0.4	MG/L
13-Mar-06	SMC1	Upstream sample site Ramona Grasslands	NH3_NO3_NO2_AUT	Nitrate	1.79	0.078	MG/L
27-Mar-06	SMC1	Upstream sample site Ramona Grasslands	ANIONS_IC_LOW	Nitrate	ND	0.1	MG/L
27-Mar-06	SMC1	Upstream sample site Ramona Grasslands	NH3_NO3_NO2_AUT	Nitrate	0.105	0.078	MG/L
3-Jan-05	SMC1	Upstream sample site Ramona Grasslands	NH3_NO3_NO2_AUT	Nitrate & Nitrite	6.99	0.3	MG/L
24-Jan-05	SMC1	Upstream sample site Ramona Grasslands	NH3_NO3_NO2_AUT	Nitrate & Nitrite	19.4	0.6	MG/L
14-Feb-05	SMC1	Upstream sample site Ramona Grasslands	NH3_NO3_NO2_AUT	Nitrate & Nitrite	19.9	0.6	MG/L
28-Feb-05	SMC1	Upstream sample site Ramona Grasslands	NH3_NO3_NO2_AUT	Nitrate & Nitrite	32.6	0.6	MG/L
15-Mar-05	SMC1	Upstream sample site Ramona Grasslands	NH3_NO3_NO2_AUT	Nitrate & Nitrite	43.2	1.2	MG/L
28-Mar-05	SMC1	Upstream sample site Ramona Grasslands	NH3_NO3_NO2_AUT	Nitrate & Nitrite	37.6	0.6	MG/L
25-Apr-05	SMC1	Upstream sample site Ramona Grasslands	NH3_NO3_NO2_AUT	Nitrate & Nitrite	> 25	0.3	MG/L
9-May-05	SMC1	Upstream sample site Ramona Grasslands	NH3_NO3_NO2_AUT	Nitrate & Nitrite	35.9	0.6	MG/L
23-May-05	SMC1	Upstream sample site Ramona Grasslands	NH3_NO3_NO2_AUT	Nitrate & Nitrite	16.6	0.6	MG/L
13-Jun-05	SMC1	Upstream sample site Ramona Grasslands	NH3_NO3_NO2_AUT	Nitrate & Nitrite	0.956	0.06	MG/L
13-Mar-06	SMC1	Upstream sample site Ramona Grasslands	NH3_NO3_NO2_AUT	Nitrate & Nitrite	1.82	0.078	MG/L
27-Mar-06	SMC1	Upstream sample site Ramona Grasslands	NH3_NO3_NO2_AUT	Nitrate & Nitrite	0.105	0.078	MG/L
3-Jan-05	SMC1	Upstream sample site Ramona Grasslands	NH3_NO3_NO2_AUT	Nitrite (NO2)	0.223	0.01	MG/L
24-Jan-05	SMC1	Upstream sample site Ramona Grasslands	NH3_NO3_NO2_AUT	Nitrite (NO2)	0.09	0.01	MG/L
14-Feb-05	SMC1	Upstream sample site Ramona Grasslands	NH3_NO3_NO2_AUT	Nitrite (NO2)	0.45	0.01	MG/L

Sample Date	Station	Source Description	Analysis	Analyte name	Value	MDL*	Units
28-Feb-05	SMC1	Upstream sample site Ramona Grasslands	NH3_NO3_NO2_AUT	Nitrite (NO2)	1	0.05	MG/L
15-Mar-05	SMC1	Upstream sample site Ramona Grasslands	NH3_NO3_NO2_AUT	Nitrite (NO2)	0.302	0.01	MG/L
28-Mar-05	SMC1	Upstream sample site Ramona Grasslands	NH3_NO3_NO2_AUT	Nitrite (NO2)	0.188	0.01	MG/L
25-Apr-05	SMC1	Upstream sample site Ramona Grasslands	NH3_NO3_NO2_AUT	Nitrite (NO2)	0.167	0.01	MG/L
9-May-05	SMC1	Upstream sample site Ramona Grasslands	NH3_NO3_NO2_AUT	Nitrite (NO2)	0.166	0.01	MG/L
23-May-05	SMC1	Upstream sample site Ramona Grasslands	NH3_NO3_NO2_AUT	Nitrite (NO2)	0.425	0.01	MG/L
13-Jun-05	SMC1	Upstream sample site Ramona Grasslands	NH3_NO3_NO2_AUT	Nitrite (NO2)	0.054	0.01	MG/L
13-Mar-06	SMC1	Upstream sample site Ramona Grasslands	NH3_NO3_NO2_AUT	Nitrite (NO2)	0.031	0.016	MG/L
27-Mar-06	SMC1	Upstream sample site Ramona Grasslands	NH3_NO3_NO2_AUT	Nitrite (NO2)	ND	0.016	MG/L
3-Jan-05	SMC1	Upstream sample site Ramona Grasslands	ANIONS_IC	Ortho phosphates	0.98	0.2	MG/L
24-Jan-05	SMC1	Upstream sample site Ramona Grasslands	ANIONS_IC_LOW	Ortho phosphates	1.21	0.1	MG/L
14-Feb-05	SMC1	Upstream sample site Ramona Grasslands	ANIONS_IC_LOW	Ortho phosphates	2.33	0.05	MG/L
28-Feb-05	SMC1	Upstream sample site Ramona Grasslands	ANIONS_IC_LOW	Ortho phosphates	1.94	0.15	MG/L
15-Mar-05	SMC1	Upstream sample site Ramona Grasslands	ANIONS_IC_LOW	Ortho phosphates	1.35	0.05	MG/L
28-Mar-05	SMC1	Upstream sample site Ramona Grasslands	ANIONS_IC_LOW	Ortho phosphates	1.62	0.05	MG/L
11-Apr-05	SMC1	Upstream sample site Ramona Grasslands	ANIONS_IC	Ortho phosphates	1.18	0.2	MG/L
25-Apr-05	SMC1	Upstream sample site Ramona Grasslands	ANIONS_IC_LOW	Ortho phosphates	1.2	0.1	MG/L
9-May-05	SMC1	Upstream sample site Ramona Grasslands	ANIONS_IC	Ortho phosphates	1.08	0.2	MG/L
23-May-05	SMC1	Upstream sample site Ramona Grasslands	ANIONS_IC	Ortho phosphates	1.13	0.2	MG/L

Sample Date	Station	Source Description	Analysis	Analyte name	Value	MDL*	Units
23-May-05	SMC1	Upstream sample site Ramona Grasslands	ANIONS_IC_LOW	Ortho phosphates	1.26	0.1	MG/L
13-Jun-05	SMC1	Upstream sample site Ramona Grasslands	ANIONS_IC	Ortho phosphates	0.675	0.2	MG/L
13-Mar-06	SMC1	Upstream sample site Ramona Grasslands	ANIONS_IC	Ortho phosphates	0.72	0.2	MG/L
27-Mar-06	SMC1	Upstream sample site Ramona Grasslands	ANIONS_IC_LOW	Ortho phosphates	0.92	0.05	MG/L
3-Jan-05	SMC1	Upstream sample site Ramona Grasslands	STREAM_PARAM	Oxidation- Reduction Potential	379		MV
24-Jan-05	SMC1	Upstream sample site Ramona Grasslands	STREAM_PARAM	Oxidation- Reduction Potential	387		MV
14-Feb-05	SMC1	Upstream sample site Ramona Grasslands	STREAM_PARAM	Oxidation- Reduction Potential	338		MV
28-Feb-05	SMC1	Upstream sample site Ramona Grasslands	STREAM_PARAM	Oxidation- Reduction Potential	335		MV
15-Mar-05	SMC1	Upstream sample site Ramona Grasslands	STREAM_PARAM	Oxidation- Reduction Potential	395		MV
28-Mar-05	SMC1	Upstream sample site Ramona Grasslands	STREAM_PARAM	Oxidation- Reduction Potential	443		MV
11-Apr-05	SMC1	Upstream sample site Ramona Grasslands	STREAM_PARAM	Oxidation- Reduction Potential	354		MV
25-Apr-05	SMC1	Upstream sample site Ramona Grasslands	STREAM_PARAM	Oxidation- Reduction Potential	424		MV
9-May-05	SMC1	Upstream sample site Ramona Grasslands	STREAM_PARAM	Oxidation- Reduction Potential	331		MV
23-May-05	SMC1	Upstream sample site Ramona Grasslands	STREAM_PARAM	Oxidation- Reduction Potential	329		MV

Sample Date	Station	Source Description	Analysis	Analyte name	Value	MDL*	Units
13-Jun-05	SMC1	Upstream sample site Ramona Grasslands	STREAM_PARAM	Oxidation- Reduction Potential	316		MV
13-Mar-06	SMC1	Upstream sample site Ramona Grasslands	STREAM_PARAM	Oxidation- Reduction Potential	364		MV
3-Jan-05	SMC1	Upstream sample site Ramona Grasslands	STREAM_PARAM	pH	7.67		PH
24-Jan-05	SMC1	Upstream sample site Ramona Grasslands	STREAM_PARAM	pH	8.13		PH
14-Feb-05	SMC1	Upstream sample site Ramona Grasslands	STREAM_PARAM	pH	7.89		PH
28-Feb-05	SMC1	Upstream sample site Ramona Grasslands	STREAM_PARAM	pH	7.88		PH
15-Mar-05	SMC1	Upstream sample site Ramona Grasslands	STREAM_PARAM	pH	7.84		PH
28-Mar-05	SMC1	Upstream sample site Ramona Grasslands	STREAM_PARAM	pH	7.95		PH
11-Apr-05	SMC1	Upstream sample site Ramona Grasslands	STREAM_PARAM	pH	8.01		PH
25-Apr-05	SMC1	Upstream sample site Ramona Grasslands	STREAM_PARAM	pH	8.26		PH
9-May-05	SMC1	Upstream sample site Ramona Grasslands	STREAM_PARAM	pH	8.28		PH
23-May-05	SMC1	Upstream sample site Ramona Grasslands	STREAM_PARAM	pH	8.38		PH
13-Jun-05	SMC1	Upstream sample site Ramona Grasslands	STREAM_PARAM	pH	8.54		PH
13-Mar-06	SMC1	Upstream sample site Ramona Grasslands	STREAM_PARAM	pH	7.87		PH
3-Jan-05	SMC1	Upstream sample site Ramona Grasslands	NIT_PHOS_UV	Phosphorus	0.421	0.084	MG/L
24-Jan-05	SMC1	Upstream sample site Ramona Grasslands	NIT_PHOS_UV	Phosphorus	0.482	0.084	MG/L
14-Feb-05	SMC1	Upstream sample site Ramona Grasslands	NIT_PHOS_UV	Phosphorus	0.914	0.084	MG/L
28-Feb-05	SMC1	Upstream sample site Ramona Grasslands	NIT_PHOS_UV	Phosphorus	0.758	0.084	MG/L

Sample Date	Station	Source Description	Analysis	Analyte name	Value	MDL*	Units
15-Mar-05	SMC1	Upstream sample site Ramona Grasslands	NIT_PHOS_UV	Phosphorus	0.696	0.084	MG/L
28-Mar-05	SMC1	Upstream sample site Ramona Grasslands	NIT_PHOS_UV	Phosphorus	0.616	0.084	MG/L
11-Apr-05	SMC1	Upstream sample site Ramona Grasslands	NIT_PHOS_UV	Phosphorus	0.584	0.084	MG/L
25-Apr-05	SMC1	Upstream sample site Ramona Grasslands	NIT_PHOS_UV	Phosphorus	0.531	0.084	MG/L
9-May-05	SMC1	Upstream sample site Ramona Grasslands	NIT_PHOS_UV	Phosphorus	0.571	0.084	MG/L
23-May-05	SMC1	Upstream sample site Ramona Grasslands	NIT_PHOS_UV	Phosphorus		0.084	MG/L
13-Jun-05	SMC1	Upstream sample site Ramona Grasslands	NIT_PHOS_UV	Phosphorus	0.511	0.084	MG/L
13-Mar-06	SMC1	Upstream sample site Ramona Grasslands	NIT_PHOS_UV	Phosphorus	0.303	0.078	MG/L
27-Mar-06	SMC1	Upstream sample site Ramona Grasslands	NIT_PHOS_UV	Phosphorus	0.341	0.078	MG/L
3-Jan-05	SMC1	Upstream sample site Ramona Grasslands	STREAM_PARAM	Stream Average Depth	0.34		FT
24-Jan-05	SMC1	Upstream sample site Ramona Grasslands	STREAM_PARAM	Stream Average Depth	1.83		FT
14-Feb-05	SMC1	Upstream sample site Ramona Grasslands	STREAM_PARAM	Stream Average Depth	0.7		FT
28-Feb-05	SMC1	Upstream sample site Ramona Grasslands	STREAM_PARAM	Stream Average Depth	0.57		FT
15-Mar-05	SMC1	Upstream sample site Ramona Grasslands	STREAM_PARAM	Stream Average Depth	0.38		FT
28-Mar-05	SMC1	Upstream sample site Ramona Grasslands	STREAM_PARAM	Stream Average Depth	0.2		FT
11-Apr-05	SMC1	Upstream sample site Ramona Grasslands	STREAM_PARAM	Stream Average Depth	0.38		FT
25-Apr-05	SMC1	Upstream sample site Ramona Grasslands	STREAM_PARAM	Stream Average Depth	0.24		FT
9-May-05	SMC1	Upstream sample site Ramona Grasslands	STREAM_PARAM	Stream Average Depth	0.19		FT
23-May-05	SMC1	Upstream sample site Ramona Grasslands	STREAM_PARAM	Stream Average Depth	0.07		FT

Sample Date	Station	Source Description	Analysis	Analyte name	Value	MDL*	Units
13-Jun-05	SMC1	Upstream sample site Ramona Grasslands	STREAM_PARAM	Stream Average Depth	0.12		FT
13-Mar-06	SMC1	Upstream sample site Ramona Grasslands	STREAM_PARAM	Stream Average Depth	0.15		FT
3-Jan-05	SMC1	Upstream sample site Ramona Grasslands	STREAM_PARAM	Stream Flow	7.85		FT3/S
24-Jan-05	SMC1	Upstream sample site Ramona Grasslands	STREAM_PARAM	Stream Flow	12.2		FT3/S
14-Feb-05	SMC1	Upstream sample site Ramona Grasslands	STREAM_PARAM	Stream Flow	21.4		FT3/S
28-Feb-05	SMC1	Upstream sample site Ramona Grasslands	STREAM_PARAM	Stream Flow	32.3		FT3/S
15-Mar-05	SMC1	Upstream sample site Ramona Grasslands	STREAM_PARAM	Stream Flow	19.4		FT3/S
28-Mar-05	SMC1	Upstream sample site Ramona Grasslands	STREAM_PARAM	Stream Flow	6.3		FT3/S
11-Apr-05	SMC1	Upstream sample site Ramona Grasslands	STREAM_PARAM	Stream Flow	4.08		FT3/S
25-Apr-05	SMC1	Upstream sample site Ramona Grasslands	STREAM_PARAM	Stream Flow	2.56		FT3/S
9-May-05	SMC1	Upstream sample site Ramona Grasslands	STREAM_PARAM	Stream Flow	2.6		FT3/S
23-May-05	SMC1	Upstream sample site Ramona Grasslands	STREAM_PARAM	Stream Flow	0.08		FT3/S
13-Jun-05	SMC1	Upstream sample site Ramona Grasslands	STREAM_PARAM	Stream Flow	0.26		FT3/S
13-Mar-06	SMC1	Upstream sample site Ramona Grasslands	STREAM_PARAM	Stream Flow	0.42		FT3/S
3-Jan-05	SMC1	Upstream sample site Ramona Grasslands	STREAM_PARAM	Stream Width	15		FT
24-Jan-05	SMC1	Upstream sample site Ramona Grasslands	STREAM_PARAM	Stream Width	8.3		FT
14-Feb-05	SMC1	Upstream sample site Ramona Grasslands	STREAM_PARAM	Stream Width	16		FT
28-Feb-05	SMC1	Upstream sample site Ramona Grasslands	STREAM_PARAM	Stream Width	29		FT
15-Mar-05	SMC1	Upstream sample site Ramona Grasslands	STREAM_PARAM	Stream Width	26		FT

Sample Date	Station	Source Description	Analysis	Analyte name	Value	MDL*	Units
28-Mar-05	SMC1	Upstream sample site Ramona Grasslands	STREAM_PARAM	Stream Width	26		FT
11-Apr-05	SMC1	Upstream sample site Ramona Grasslands	STREAM_PARAM	Stream Width	7		FT
25-Apr-05	SMC1	Upstream sample site Ramona Grasslands	STREAM_PARAM	Stream Width	10.2		FT
9-May-05	SMC1	Upstream sample site Ramona Grasslands	STREAM_PARAM	Stream Width	12		FT
23-May-05	SMC1	Upstream sample site Ramona Grasslands	STREAM_PARAM	Stream Width	3.2		FT
13-Jun-05	SMC1	Upstream sample site Ramona Grasslands	STREAM_PARAM	Stream Width	3.4		FT
13-Mar-06	SMC1	Upstream sample site Ramona Grasslands	STREAM_PARAM	Stream Width	4.2		FT
3-Jan-05	SMC1	Upstream sample site Ramona Grasslands	ANIONS_IC	Sulfate	31.8	0.5	MG/L
28-Feb-05	SMC1	Upstream sample site Ramona Grasslands	ANIONS_IC	Sulfate	77.3	0.5	MG/L
11-Apr-05	SMC1	Upstream sample site Ramona Grasslands	ANIONS_IC	Sulfate	126	0.5	MG/L
25-Apr-05	SMC1	Upstream sample site Ramona Grasslands	ANIONS_IC	Sulfate	128	0.5	MG/L
9-May-05	SMC1	Upstream sample site Ramona Grasslands	ANIONS_IC	Sulfate	124	0.5	MG/L
23-May-05	SMC1	Upstream sample site Ramona Grasslands	ANIONS_IC	Sulfate	142	0.5	MG/L
13-Jun-05	SMC1	Upstream sample site Ramona Grasslands	ANIONS_IC	Sulfate	154	0.5	MG/L
13-Mar-06	SMC1	Upstream sample site Ramona Grasslands	ANIONS_IC	Sulfate	83.4	0.5	MG/L
27-Mar-06	SMC1	Upstream sample site Ramona Grasslands	ANIONS_IC_LOW	Sulfate	104	0.125	MG/L
3-Jan-05	SMC1	Upstream sample site Ramona Grasslands	STREAM_PARAM	Temperature degrees C	9.03		DEGREE_C
24-Jan-05	SMC1	Upstream sample site Ramona Grasslands	STREAM_PARAM	Temperature degrees C	13.2		DEGREE_C
14-Feb-05	SMC1	Upstream sample site Ramona Grasslands	STREAM_PARAM	Temperature degrees C	12.2		DEGREE_C

Sample Date	Station	Source Description	Analysis	Analyte name	Value	MDL*	Units
28-Feb-05	SMC1	Upstream sample site Ramona Grasslands	STREAM_PARAM	Temperature degrees C	11.2		DEGREE_C
15-Mar-05	SMC1	Upstream sample site Ramona Grasslands	STREAM_PARAM	Temperature degrees C	8.57		DEGREE_C
28-Mar-05	SMC1	Upstream sample site Ramona Grasslands	STREAM_PARAM	Temperature degrees C	11.5		DEGREE_C
11-Apr-05	SMC1	Upstream sample site Ramona Grasslands	STREAM_PARAM	Temperature degrees C	9.59		DEGREE_C
25-Apr-05	SMC1	Upstream sample site Ramona Grasslands	STREAM_PARAM	Temperature degrees C	13.5		DEGREE_C
9-May-05	SMC1	Upstream sample site Ramona Grasslands	STREAM_PARAM	Temperature degrees C	15.2		DEGREE_C
23-May-05	SMC1	Upstream sample site Ramona Grasslands	STREAM_PARAM	Temperature degrees C	19.4		DEGREE_C
13-Jun-05	SMC1	Upstream sample site Ramona Grasslands	STREAM_PARAM	Temperature degrees C	21.7		DEGREE_C
13-Mar-06	SMC1	Upstream sample site Ramona Grasslands	STREAM_PARAM	Temperature degrees C	6.16		DEGREE_C
3-Jan-05	SMC1	Upstream sample site Ramona Grasslands	QT_CST	Total Coliform	> 240000		/100 ML
24-Jan-05	SMC1	Upstream sample site Ramona Grasslands	QT_CST	Total Coliform	3100		/100 ML
14-Feb-05	SMC1	Upstream sample site Ramona Grasslands	QT_CST	Total Coliform	20000		/100 ML
28-Feb-05	SMC1	Upstream sample site Ramona Grasslands	QT_CST	Total Coliform	16000		/100 ML
15-Mar-05	SMC1	Upstream sample site Ramona Grasslands	QT_CST	Total Coliform	22000		/100 ML
28-Mar-05	SMC1	Upstream sample site Ramona Grasslands	QT_CST	Total Coliform	18000		/100 ML
11-Apr-05	SMC1	Upstream sample site Ramona Grasslands	QT_CST	Total Coliform	9300		/100 ML
25-Apr-05	SMC1	Upstream sample site Ramona Grasslands	QT_CST	Total Coliform	12000		/100 ML
9-May-05	SMC1	Upstream sample site Ramona Grasslands	QT_CST	Total Coliform	13000		/100 ML
23-May-05	SMC1	Upstream sample site Ramona Grasslands	QT_CST	Total Coliform	100000		/100 ML

Sample Date	Station	Source Description	Analysis	Analyte name	Value	MDL*	Units
13-Jun-05	SMC1	Upstream sample site Ramona Grasslands	QT_CST	Total Coliform	77000		/100 ML
13-Mar-06	SMC1	Upstream sample site Ramona Grasslands	QT_CST	Total Coliform	10000		/100 ML
27-Mar-06	SMC1	Upstream sample site Ramona Grasslands	QT_CST	Total Coliform	3900		/100 ML
3-Jan-05	SMC1	Upstream sample site Ramona Grasslands	SOLIDS_TDS	Total Dissolved Solids	360	20	MG/L
24-Jan-05	SMC1	Upstream sample site Ramona Grasslands	SOLIDS_TDS	Total Dissolved Solids	982	10	MG/L
14-Feb-05	SMC1	Upstream sample site Ramona Grasslands	SOLIDS_TDS	Total Dissolved Solids	743	10	MG/L
28-Feb-05	SMC1	Upstream sample site Ramona Grasslands	SOLIDS_TDS	Total Dissolved Solids	564	10	MG/L
15-Mar-05	SMC1	Upstream sample site Ramona Grasslands	SOLIDS_TDS	Total Dissolved Solids	706	10	MG/L
28-Mar-05	SMC1	Upstream sample site Ramona Grasslands	SOLIDS_TDS	Total Dissolved Solids	703	10	MG/L
11-Apr-05	SMC1	Upstream sample site Ramona Grasslands	SOLIDS_TDS	Total Dissolved Solids	764	10	MG/L
25-Apr-05	SMC1	Upstream sample site Ramona Grasslands	SOLIDS_TDS	Total Dissolved Solids	811	10	MG/L
9-May-05	SMC1	Upstream sample site Ramona Grasslands	SOLIDS_TDS	Total Dissolved Solids	772	10	MG/L
23-May-05	SMC1	Upstream sample site Ramona Grasslands	SOLIDS_TDS	Total Dissolved Solids	892	10	MG/L
13-Jun-05	SMC1	Upstream sample site Ramona Grasslands	SOLIDS_TDS	Total Dissolved Solids	968	10	MG/L
13-Mar-06	SMC1	Upstream sample site Ramona Grasslands	SOLIDS_TDS	Total Dissolved Solids	608	10	MG/L
27-Mar-06	SMC1	Upstream sample site Ramona Grasslands	SOLIDS_TDS	Total Dissolved Solids	796	10	MG/L
3-Jan-05	SMC1	Upstream sample site Ramona Grasslands	NIT_PHOS_UV	Total Nitrogen	2.15	0.22	MG/L
24-Jan-05	SMC1	Upstream sample site Ramona Grasslands	NIT_PHOS_UV	Total Nitrogen	5.51	0.22	MG/L
14-Feb-05	SMC1	Upstream sample site Ramona Grasslands	NIT_PHOS_UV	Total Nitrogen	6.76	0.22	MG/L

Sample Date	Station	Source Description	Analysis	Analyte name	Value	MDL*	Units
28-Feb-05	SMC1	Upstream sample site Ramona Grasslands	NIT_PHOS_UV	Total Nitrogen	> 10	0.22	MG/L
15-Mar-05	SMC1	Upstream sample site Ramona Grasslands	NIT_PHOS_UV	Total Nitrogen	> 10	0.22	MG/L
28-Mar-05	SMC1	Upstream sample site Ramona Grasslands	NIT_PHOS_UV	Total Nitrogen	9.85	0.22	MG/L
11-Apr-05	SMC1	Upstream sample site Ramona Grasslands	NIT_PHOS_UV	Total Nitrogen	9.92	0.22	MG/L
25-Apr-05	SMC1	Upstream sample site Ramona Grasslands	NIT_PHOS_UV	Total Nitrogen	10.8	0.44	MG/L
9-May-05	SMC1	Upstream sample site Ramona Grasslands	NIT_PHOS_UV	Total Nitrogen	9.2	0.22	MG/L
23-May-05	SMC1	Upstream sample site Ramona Grasslands	NIT_PHOS_UV	Total Nitrogen		0.22	MG/L
13-Jun-05	SMC1	Upstream sample site Ramona Grasslands	NIT_PHOS_UV	Total Nitrogen	0.642	0.22	MG/L
13-Mar-06	SMC1	Upstream sample site Ramona Grasslands	NIT_PHOS_UV	Total Nitrogen	1.03	0.156	MG/L
27-Mar-06	SMC1	Upstream sample site Ramona Grasslands	NIT_PHOS_UV	Total Nitrogen	0.302	0.156	MG/L
3-Jan-05	SMC1	Upstream sample site Ramona Grasslands	SOLIDS_TSS	Total Suspended Solids (TSS)	252		MG/L
24-Jan-05	SMC1	Upstream sample site Ramona Grasslands	SOLIDS_TSS	Total Suspended Solids (TSS)	< 1	1	MG/L
14-Feb-05	SMC1	Upstream sample site Ramona Grasslands	SOLIDS_TSS	Total Suspended Solids (TSS)	15.1	1	MG/L
28-Feb-05	SMC1	Upstream sample site Ramona Grasslands	SOLIDS_TSS	Total Suspended Solids (TSS)	73		MG/L
15-Mar-05	SMC1	Upstream sample site Ramona Grasslands	SOLIDS_TSS	Total Suspended Solids (TSS)	15.5	1	MG/L
28-Mar-05	SMC1	Upstream sample site Ramona Grasslands	SOLIDS_TSS	Total Suspended Solids (TSS)	22.2		MG/L
11-Apr-05	SMC1	Upstream sample site Ramona Grasslands	SOLIDS_TSS	Total Suspended Solids (TSS)	5.7		MG/L
25-Apr-05	SMC1	Upstream sample site Ramona Grasslands	SOLIDS_TSS	Total Suspended Solids (TSS)	3.8		MG/L
9-May-05	SMC1	Upstream sample site Ramona Grasslands	SOLIDS_TSS	Total Suspended Solids (TSS)	6.7		MG/L

Sample Date	Station	Source Description	Analysis	Analyte name	Value	MDL*	Units
23-May-05	SMC1	Upstream sample site Ramona Grasslands	SOLIDS_TSS	Total Suspended Solids (TSS)	4.2		MG/L
13-Jun-05	SMC1	Upstream sample site Ramona Grasslands	SOLIDS_TSS	Total Suspended Solids (TSS)	5.1		MG/L
13-Mar-06	SMC1	Upstream sample site Ramona Grasslands	SOLIDS_TSS	Total Suspended Solids (TSS)	5.5		MG/L
27-Mar-06	SMC1	Upstream sample site Ramona Grasslands	SOLIDS_TSS	Total Suspended Solids (TSS)	7.3	1	MG/L
3-Jan-05	SMC2	Downstream sample site Ramona Grasslands	NH3_NO3_NO2_AUT	Ammonia-N	0.198	0.04	MG/L
24-Jan-05	SMC2	Downstream sample site Ramona Grasslands	NH3_NO3_NO2_AUT	Ammonia-N	0.122	0.04	MG/L
14-Feb-05	SMC2	Downstream sample site Ramona Grasslands	NH3_NO3_NO2_AUT	Ammonia-N	0.872	0.04	MG/L
28-Feb-05	SMC2	Downstream sample site Ramona Grasslands	NH3_NO3_NO2_AUT	Ammonia-N	0.605	0.04	MG/L
15-Mar-05	SMC2	Downstream sample site Ramona Grasslands	NH3_NO3_NO2_AUT	Ammonia-N	0.079	0.04	MG/L
28-Mar-05	SMC2	Downstream sample site Ramona Grasslands	NH3_NO3_NO2_AUT	Ammonia-N	0.095	0.04	MG/L
11-Apr-05	SMC2	Downstream sample site Ramona Grasslands	NH3_NO3_NO2_AUT	Ammonia-N	ND	0.04	MG/L
25-Apr-05	SMC2	Downstream sample site Ramona Grasslands	NH3_NO3_NO2_AUT	Ammonia-N	0.048	0.04	MG/L
9-May-05	SMC2	Downstream sample site Ramona Grasslands	NH3_NO3_NO2_AUT	Ammonia-N	ND	0.04	MG/L
23-May-05	SMC2	Downstream sample site Ramona Grasslands	NH3_NO3_NO2_AUT	Ammonia-N	0.05	0.04	MG/L
13-Mar-06	SMC2	Downstream sample site Ramona Grasslands	NH3_NO3_NO2_AUT	Ammonia-N	0.066	0.031	MG/L
3-Jan-05	SMC2	Downstream sample site Ramona Grasslands	ANIONS_IC	Bromide	0.231	0.1	MG/L
28-Feb-05	SMC2	Downstream sample site Ramona Grasslands	ANIONS_IC	Bromide	0.254	0.1	MG/L
11-Apr-05	SMC2	Downstream sample site Ramona Grasslands	ANIONS_IC	Bromide	0.428	0.1	MG/L
25-Apr-05	SMC2	Downstream sample site Ramona Grasslands	ANIONS_IC	Bromide	0.407	0.1	MG/L

Sample Date	Station	Source Description	Analysis	Analyte name	Value	MDL*	Units
9-May-05	SMC2	Downstream sample site Ramona Grasslands	ANIONS_IC	Bromide	0.433	0.1	MG/L
23-May-05	SMC2	Downstream sample site Ramona Grasslands	ANIONS_IC	Bromide	1.86	0.1	MG/L
13-Mar-06	SMC2	Downstream sample site Ramona Grasslands	ANIONS_IC	Bromide	1.21	0.1	MG/L
3-Jan-05	SMC2	Downstream sample site Ramona Grasslands	ANIONS_IC	Chloride	94.1	0.5	MG/L
28-Feb-05	SMC2	Downstream sample site Ramona Grasslands	ANIONS_IC	Chloride	109	1	MG/L
11-Apr-05	SMC2	Downstream sample site Ramona Grasslands	ANIONS_IC	Chloride	165	1	MG/L
25-Apr-05	SMC2	Downstream sample site Ramona Grasslands	ANIONS_IC	Chloride	167	1	MG/L
9-May-05	SMC2	Downstream sample site Ramona Grasslands	ANIONS_IC	Chloride	159	1	MG/L
13-Mar-06	SMC2	Downstream sample site Ramona Grasslands	ANIONS_IC	Chloride	215	0.5	MG/L
3-Jan-05	SMC2	Downstream sample site Ramona Grasslands	STREAM_PARAM	Conductivity	568		US/CM
24-Jan-05	SMC2	Downstream sample site Ramona Grasslands	STREAM_PARAM	Conductivity	1430		US/CM
14-Feb-05	SMC2	Downstream sample site Ramona Grasslands	STREAM_PARAM	Conductivity	963		US/CM
28-Feb-05	SMC2	Downstream sample site Ramona Grasslands	STREAM_PARAM	Conductivity	938		US/CM
15-Mar-05	SMC2	Downstream sample site Ramona Grasslands	STREAM_PARAM	Conductivity	1210		US/CM
28-Mar-05	SMC2	Downstream sample site Ramona Grasslands	STREAM_PARAM	Conductivity	1220		US/CM
11-Apr-05	SMC2	Downstream sample site Ramona Grasslands	STREAM_PARAM	Conductivity	1290		US/CM
25-Apr-05	SMC2	Downstream sample site Ramona Grasslands	STREAM_PARAM	Conductivity	1240		US/CM
9-May-05	SMC2	Downstream sample site Ramona Grasslands	STREAM_PARAM	Conductivity	1230		US/CM
23-May-05	SMC2	Downstream sample site Ramona Grasslands	STREAM_PARAM	Conductivity	2510		US/CM

Sample Date	Station	Source Description	Analysis	Analyte name	Value	MDL*	Units
13-Mar-06	SMC2	Downstream sample site Ramona Grasslands	STREAM_PARAM	Conductivity	1160		US/CM
3-Jan-05	SMC2	Downstream sample site Ramona Grasslands	STREAM_PARAM	Dissolved Oxygen	8.39		MG/L
24-Jan-05	SMC2	Downstream sample site Ramona Grasslands	STREAM_PARAM	Dissolved Oxygen	10		MG/L
14-Feb-05	SMC2	Downstream sample site Ramona Grasslands	STREAM_PARAM	Dissolved Oxygen	8.43		MG/L
28-Feb-05	SMC2	Downstream sample site Ramona Grasslands	STREAM_PARAM	Dissolved Oxygen	9.7		MG/L
15-Mar-05	SMC2	Downstream sample site Ramona Grasslands	STREAM_PARAM	Dissolved Oxygen	10.5		MG/L
28-Mar-05	SMC2	Downstream sample site Ramona Grasslands	STREAM_PARAM	Dissolved Oxygen	9.71		MG/L
11-Apr-05	SMC2	Downstream sample site Ramona Grasslands	STREAM_PARAM	Dissolved Oxygen	10.9		MG/L
25-Apr-05	SMC2	Downstream sample site Ramona Grasslands	STREAM_PARAM	Dissolved Oxygen	11.1		MG/L
9-May-05	SMC2	Downstream sample site Ramona Grasslands	STREAM_PARAM	Dissolved Oxygen	8.72		MG/L
23-May-05	SMC2	Downstream sample site Ramona Grasslands	STREAM_PARAM	Dissolved Oxygen	9.02		MG/L
13-Mar-06	SMC2	Downstream sample site Ramona Grasslands	STREAM_PARAM	Dissolved Oxygen	10.4		MG/L
3-Jan-05	SMC2	Downstream sample site Ramona Grasslands	QT_CST	Enterococcus	130		/100 ML
24-Jan-05	SMC2	Downstream sample site Ramona Grasslands	QT_CST	Enterococcus	230		/100 ML
14-Feb-05	SMC2	Downstream sample site Ramona Grasslands	QT_CST	Enterococcus	1600		/100 ML
28-Feb-05	SMC2	Downstream sample site Ramona Grasslands	QT_CST	Enterococcus	120		/100 ML
15-Mar-05	SMC2	Downstream sample site Ramona Grasslands	QT_CST	Enterococcus	93		/100 ML
28-Mar-05	SMC2	Downstream sample site Ramona Grasslands	QT_CST	Enterococcus	75		/100 ML
11-Apr-05	SMC2	Downstream sample site Ramona Grasslands	QT_CST	Enterococcus	36		/100 ML

Sample Date	Station	Source Description	Analysis	Analyte name	Value	MDL*	Units
25-Apr-05	SMC2	Downstream sample site Ramona Grasslands	QT_CST	Enterococcus	16		/100 ML
9-May-05	SMC2	Downstream sample site Ramona Grasslands	QT_CST	Enterococcus	120		/100 ML
23-May-05	SMC2	Downstream sample site Ramona Grasslands	QT_CST	Enterococcus	820		/100 ML
13-Mar-06	SMC2	Downstream sample site Ramona Grasslands	QT_CST	Enterococcus	1100		/100 ML
3-Jan-05	SMC2	Downstream sample site Ramona Grasslands	STREAM_PARAM	Equipment Number	39300		.
24-Jan-05	SMC2	Downstream sample site Ramona Grasslands	STREAM_PARAM	Equipment Number	39300		.
14-Feb-05	SMC2	Downstream sample site Ramona Grasslands	STREAM_PARAM	Equipment Number	39300		.
28-Feb-05	SMC2	Downstream sample site Ramona Grasslands	STREAM_PARAM	Equipment Number	39300		.
15-Mar-05	SMC2	Downstream sample site Ramona Grasslands	STREAM_PARAM	Equipment Number	39300		.
28-Mar-05	SMC2	Downstream sample site Ramona Grasslands	STREAM_PARAM	Equipment Number	39300		.
11-Apr-05	SMC2	Downstream sample site Ramona Grasslands	STREAM_PARAM	Equipment Number	39300		.
25-Apr-05	SMC2	Downstream sample site Ramona Grasslands	STREAM_PARAM	Equipment Number	39300		.
9-May-05	SMC2	Downstream sample site Ramona Grasslands	STREAM_PARAM	Equipment Number	39300		.
23-May-05	SMC2	Downstream sample site Ramona Grasslands	STREAM_PARAM	Equipment Number	39300		.
3-Jan-05	SMC2	Downstream sample site Ramona Grasslands	QT_CST	Escherichia Coli	1300		/100 ML
24-Jan-05	SMC2	Downstream sample site Ramona Grasslands	QT_CST	Escherichia Coli	310		/100 ML
14-Feb-05	SMC2	Downstream sample site Ramona Grasslands	QT_CST	Escherichia Coli	5700		/100 ML
28-Feb-05	SMC2	Downstream sample site Ramona Grasslands	QT_CST	Escherichia Coli	200		/100 ML
15-Mar-05	SMC2	Downstream sample site Ramona Grasslands	QT_CST	Escherichia Coli	740		/100 ML

Sample Date	Station	Source Description	Analysis	Analyte name	Value	MDL*	Units
28-Mar-05	SMC2	Downstream sample site Ramona Grasslands	QT_CST	Escherichia Coli	1300		/100 ML
11-Apr-05	SMC2	Downstream sample site Ramona Grasslands	QT_CST	Escherichia Coli	630		/100 ML
25-Apr-05	SMC2	Downstream sample site Ramona Grasslands	QT_CST	Escherichia Coli	1500		/100 ML
9-May-05	SMC2	Downstream sample site Ramona Grasslands	QT_CST	Escherichia Coli	620		/100 ML
23-May-05	SMC2	Downstream sample site Ramona Grasslands	QT_CST	Escherichia Coli	< 100		/100 ML
13-Mar-06	SMC2	Downstream sample site Ramona Grasslands	QT_CST	Escherichia Coli	630		/100 ML
3-Jan-05	SMC2	Downstream sample site Ramona Grasslands	NH3_NO3_NO2_AUT	Nitrate	3.77	0.06	MG/L
24-Jan-05	SMC2	Downstream sample site Ramona Grasslands	NH3_NO3_NO2_AUT	Nitrate	20.3	0.6	MG/L
14-Feb-05	SMC2	Downstream sample site Ramona Grasslands	NH3_NO3_NO2_AUT	Nitrate	11.7	0.6	MG/L
28-Feb-05	SMC2	Downstream sample site Ramona Grasslands	NH3_NO3_NO2_AUT	Nitrate	26.7	0.6	MG/L
15-Mar-05	SMC2	Downstream sample site Ramona Grasslands	NH3_NO3_NO2_AUT	Nitrate	33.1	1.2	MG/L
28-Mar-05	SMC2	Downstream sample site Ramona Grasslands	NH3_NO3_NO2_AUT	Nitrate	29.3	0.6	MG/L
11-Apr-05	SMC2	Downstream sample site Ramona Grasslands	ANIONS_IC	Nitrate	35.9	0.4	MG/L
25-Apr-05	SMC2	Downstream sample site Ramona Grasslands	NH3_NO3_NO2_AUT	Nitrate	> 25	0.3	MG/L
9-May-05	SMC2	Downstream sample site Ramona Grasslands	ANIONS_IC	Nitrate	26.8	0.2	MG/L
9-May-05	SMC2	Downstream sample site Ramona Grasslands	NH3_NO3_NO2_AUT	Nitrate	24.9	0.3	MG/L
23-May-05	SMC2	Downstream sample site Ramona Grasslands	NH3_NO3_NO2_AUT	Nitrate	ND	0.06	MG/L
13-Mar-06	SMC2	Downstream sample site Ramona Grasslands	ANIONS_IC	Nitrate	3.95	0.4	MG/L
13-Mar-06	SMC2	Downstream sample site Ramona Grasslands	NH3_NO3_NO2_AUT	Nitrate	3.08	0.078	MG/L

Sample Date	Station	Source Description	Analysis	Analyte name	Value	MDL*	Units
3-Jan-05	SMC2	Downstream sample site Ramona Grasslands	NH3_NO3_NO2_AUT	Nitrate & Nitrite	3.91	0.06	MG/L
24-Jan-05	SMC2	Downstream sample site Ramona Grasslands	NH3_NO3_NO2_AUT	Nitrate & Nitrite	20.6	0.6	MG/L
14-Feb-05	SMC2	Downstream sample site Ramona Grasslands	NH3_NO3_NO2_AUT	Nitrate & Nitrite	12.2	0.6	MG/L
28-Feb-05	SMC2	Downstream sample site Ramona Grasslands	NH3_NO3_NO2_AUT	Nitrate & Nitrite	27.7	0.6	MG/L
15-Mar-05	SMC2	Downstream sample site Ramona Grasslands	NH3_NO3_NO2_AUT	Nitrate & Nitrite	33.7	1.2	MG/L
28-Mar-05	SMC2	Downstream sample site Ramona Grasslands	NH3_NO3_NO2_AUT	Nitrate & Nitrite	29.7	0.6	MG/L
25-Apr-05	SMC2	Downstream sample site Ramona Grasslands	NH3_NO3_NO2_AUT	Nitrate & Nitrite	> 25	0.3	MG/L
9-May-05	SMC2	Downstream sample site Ramona Grasslands	NH3_NO3_NO2_AUT	Nitrate & Nitrite	25	0.6	MG/L
23-May-05	SMC2	Downstream sample site Ramona Grasslands	NH3_NO3_NO2_AUT	Nitrate & Nitrite	ND	0.06	MG/L
13-Mar-06	SMC2	Downstream sample site Ramona Grasslands	NH3_NO3_NO2_AUT	Nitrate & Nitrite	3.15	0.078	MG/L
3-Jan-05	SMC2	Downstream sample site Ramona Grasslands	NH3_NO3_NO2_AUT	Nitrite (NO2)	0.14	0.01	MG/L
24-Jan-05	SMC2	Downstream sample site Ramona Grasslands	NH3_NO3_NO2_AUT	Nitrite (NO2)	0.365	0.01	MG/L
14-Feb-05	SMC2	Downstream sample site Ramona Grasslands	NH3_NO3_NO2_AUT	Nitrite (NO2)	0.46	0.01	MG/L
28-Feb-05	SMC2	Downstream sample site Ramona Grasslands	NH3_NO3_NO2_AUT	Nitrite (NO2)	1.08	0.05	MG/L
15-Mar-05	SMC2	Downstream sample site Ramona Grasslands	NH3_NO3_NO2_AUT	Nitrite (NO2)	0.431	0.01	MG/L
28-Mar-05	SMC2	Downstream sample site Ramona Grasslands	NH3_NO3_NO2_AUT	Nitrite (NO2)	0.363	0.01	MG/L
25-Apr-05	SMC2	Downstream sample site Ramona Grasslands	NH3_NO3_NO2_AUT	Nitrite (NO2)	0.19	0.01	MG/L
9-May-05	SMC2	Downstream sample site Ramona Grasslands	NH3_NO3_NO2_AUT	Nitrite (NO2)	0.174	0.01	MG/L
23-May-05	SMC2	Downstream sample site Ramona Grasslands	NH3_NO3_NO2_AUT	Nitrite (NO2)	ND	0.01	MG/L

Sample Date	Station	Source Description	Analysis	Analyte name	Value	MDL*	Units
13-Mar-06	SMC2	Downstream sample site Ramona Grasslands	NH3_NO3_NO2_AUT	Nitrite (NO2)	0.077	0.016	MG/L
3-Jan-05	SMC2	Downstream sample site Ramona Grasslands	ANIONS_IC	Ortho phosphates	1.38	0.2	MG/L
24-Jan-05	SMC2	Downstream sample site Ramona Grasslands	ANIONS_IC_LOW	Ortho phosphates	1.72	0.2	MG/L
14-Feb-05	SMC2	Downstream sample site Ramona Grasslands	ANIONS_IC_LOW	Ortho phosphates	3.49	0.05	MG/L
28-Feb-05	SMC2	Downstream sample site Ramona Grasslands	ANIONS_IC_LOW	Ortho phosphates	2.66	0.15	MG/L
15-Mar-05	SMC2	Downstream sample site Ramona Grasslands	ANIONS_IC_LOW	Ortho phosphates	1.57	0.05	MG/L
28-Mar-05	SMC2	Downstream sample site Ramona Grasslands	ANIONS_IC_LOW	Ortho phosphates	1.85	0.05	MG/L
11-Apr-05	SMC2	Downstream sample site Ramona Grasslands	ANIONS_IC	Ortho phosphates	1.38	0.2	MG/L
25-Apr-05	SMC2	Downstream sample site Ramona Grasslands	ANIONS_IC_LOW	Ortho phosphates	1.35	0.1	MG/L
9-May-05	SMC2	Downstream sample site Ramona Grasslands	ANIONS_IC	Ortho phosphates	1.24	0.2	MG/L
23-May-05	SMC2	Downstream sample site Ramona Grasslands	ANIONS_IC	Ortho phosphates	1.16	0.2	MG/L
23-May-05	SMC2	Downstream sample site Ramona Grasslands	ANIONS_IC_LOW	Ortho phosphates	1.22	0.1	MG/L
13-Mar-06	SMC2	Downstream sample site Ramona Grasslands	ANIONS_IC	Ortho phosphates	1.66	0.2	MG/L
3-Jan-05	SMC2	Downstream sample site Ramona Grasslands	STREAM_PARAM	Oxidation- Reduction Potential	394		MV
24-Jan-05	SMC2	Downstream sample site Ramona Grasslands	STREAM_PARAM	Oxidation- Reduction Potential	363		MV
14-Feb-05	SMC2	Downstream sample site Ramona Grasslands	STREAM_PARAM	Oxidation- Reduction Potential	358		MV
28-Feb-05	SMC2	Downstream sample site Ramona Grasslands	STREAM_PARAM	Oxidation- Reduction Potential	361		MV

Sample Date	Station	Source Description	Analysis	Analyte name	Value	MDL*	Units
15-Mar-05	SMC2	Downstream sample site Ramona Grasslands	STREAM_PARAM	Oxidation- Reduction Potential	358		MV
28-Mar-05	SMC2	Downstream sample site Ramona Grasslands	STREAM_PARAM	Oxidation- Reduction Potential	430		MV
11-Apr-05	SMC2	Downstream sample site Ramona Grasslands	STREAM_PARAM	Oxidation- Reduction Potential	332		MV
25-Apr-05	SMC2	Downstream sample site Ramona Grasslands	STREAM_PARAM	Oxidation- Reduction Potential	437		MV
9-May-05	SMC2	Downstream sample site Ramona Grasslands	STREAM_PARAM	Oxidation- Reduction Potential	321		MV
23-May-05	SMC2	Downstream sample site Ramona Grasslands	STREAM_PARAM	Oxidation- Reduction Potential	304		MV
13-Mar-06	SMC2	Downstream sample site Ramona Grasslands	STREAM_PARAM	Oxidation- Reduction Potential	317		MV
3-Jan-05	SMC2	Downstream sample site Ramona Grasslands	STREAM_PARAM	pH	7.68		PH
24-Jan-05	SMC2	Downstream sample site Ramona Grasslands	STREAM_PARAM	pH	7.85		PH
14-Feb-05	SMC2	Downstream sample site Ramona Grasslands	STREAM_PARAM	pH	7.64		PH
28-Feb-05	SMC2	Downstream sample site Ramona Grasslands	STREAM_PARAM	pH	7.71		PH
15-Mar-05	SMC2	Downstream sample site Ramona Grasslands	STREAM_PARAM	pH	8		PH
28-Mar-05	SMC2	Downstream sample site Ramona Grasslands	STREAM_PARAM	pH	8.05		PH
11-Apr-05	SMC2	Downstream sample site Ramona Grasslands	STREAM_PARAM	pH	8.05		PH
25-Apr-05	SMC2	Downstream sample site Ramona Grasslands	STREAM_PARAM	pH	8.54		PH
9-May-05	SMC2	Downstream sample site Ramona Grasslands	STREAM_PARAM	pH	8.31		PH

Sample Date	Station	Source Description	Analysis	Analyte name	Value	MDL*	Units
23-May-05	SMC2	Downstream sample site Ramona Grasslands	STREAM_PARAM	pH	8.24		PH
13-Mar-06	SMC2	Downstream sample site Ramona Grasslands	STREAM_PARAM	pH	7.96		PH
3-Jan-05	SMC2	Downstream sample site Ramona Grasslands	NIT_PHOS_UV	Phosphorus	0.603	0.084	MG/L
24-Jan-05	SMC2	Downstream sample site Ramona Grasslands	NIT_PHOS_UV	Phosphorus	0.694	0.084	MG/L
14-Feb-05	SMC2	Downstream sample site Ramona Grasslands	NIT_PHOS_UV	Phosphorus	1.31	0.084	MG/L
28-Feb-05	SMC2	Downstream sample site Ramona Grasslands	NIT_PHOS_UV	Phosphorus	0.981	0.084	MG/L
15-Mar-05	SMC2	Downstream sample site Ramona Grasslands	NIT_PHOS_UV	Phosphorus	0.85	0.084	MG/L
28-Mar-05	SMC2	Downstream sample site Ramona Grasslands	NIT_PHOS_UV	Phosphorus	0.811	0.084	MG/L
11-Apr-05	SMC2	Downstream sample site Ramona Grasslands	NIT_PHOS_UV	Phosphorus	0.612	0.084	MG/L
25-Apr-05	SMC2	Downstream sample site Ramona Grasslands	NIT_PHOS_UV	Phosphorus	0.542	0.084	MG/L
9-May-05	SMC2	Downstream sample site Ramona Grasslands	NIT_PHOS_UV	Phosphorus	0.59	0.084	MG/L
23-May-05	SMC2	Downstream sample site Ramona Grasslands	NIT_PHOS_UV	Phosphorus		0.084	MG/L
13-Mar-06	SMC2	Downstream sample site Ramona Grasslands	NIT_PHOS_UV	Phosphorus	0.547	0.078	MG/L
3-Jan-05	SMC2	Downstream sample site Ramona Grasslands	STREAM_PARAM	Stream Average Depth	0.5		FT
24-Jan-05	SMC2	Downstream sample site Ramona Grasslands	STREAM_PARAM	Stream Average Depth	0.33		FT
14-Feb-05	SMC2	Downstream sample site Ramona Grasslands	STREAM_PARAM	Stream Average Depth	0.28		FT
28-Feb-05	SMC2	Downstream sample site Ramona Grasslands	STREAM_PARAM	Stream Average Depth	0.97		FT
15-Mar-05	SMC2	Downstream sample site Ramona Grasslands	STREAM_PARAM	Stream Average Depth	0.36		FT
28-Mar-05	SMC2	Downstream sample site Ramona Grasslands	STREAM_PARAM	Stream Average Depth	0.45		FT

Sample Date	Station	Source Description	Analysis	Analyte name	Value	MDL*	Units
11-Apr-05	SMC2	Downstream sample site Ramona Grasslands	STREAM_PARAM	Stream Average Depth	0.24		FT
25-Apr-05	SMC2	Downstream sample site Ramona Grasslands	STREAM_PARAM	Stream Average Depth	0.18		FT
9-May-05	SMC2	Downstream sample site Ramona Grasslands	STREAM_PARAM	Stream Average Depth	0.32		FT
23-May-05	SMC2	Downstream sample site Ramona Grasslands	STREAM_PARAM	Stream Average Depth	0.08		FT
13-Mar-06	SMC2	Downstream sample site Ramona Grasslands	STREAM_PARAM	Stream Average Depth	0.23		FT
3-Jan-05	SMC2	Downstream sample site Ramona Grasslands	STREAM_PARAM	Stream Flow	7.77		FT3/S
24-Jan-05	SMC2	Downstream sample site Ramona Grasslands	STREAM_PARAM	Stream Flow	4.4		FT3/S
14-Feb-05	SMC2	Downstream sample site Ramona Grasslands	STREAM_PARAM	Stream Flow	6.82		FT3/S
28-Feb-05	SMC2	Downstream sample site Ramona Grasslands	STREAM_PARAM	Stream Flow	44.9		FT3/S
15-Mar-05	SMC2	Downstream sample site Ramona Grasslands	STREAM_PARAM	Stream Flow	10.4		FT3/S
28-Mar-05	SMC2	Downstream sample site Ramona Grasslands	STREAM_PARAM	Stream Flow	6.83		FT3/S
11-Apr-05	SMC2	Downstream sample site Ramona Grasslands	STREAM_PARAM	Stream Flow	3.49		FT3/S
25-Apr-05	SMC2	Downstream sample site Ramona Grasslands	STREAM_PARAM	Stream Flow	3.72		FT3/S
9-May-05	SMC2	Downstream sample site Ramona Grasslands	STREAM_PARAM	Stream Flow	3.5		FT3/S
23-May-05	SMC2	Downstream sample site Ramona Grasslands	STREAM_PARAM	Stream Flow	0.05		FT3/S
13-Mar-06	SMC2	Downstream sample site Ramona Grasslands	STREAM_PARAM	Stream Flow	0.26		FT3/S
3-Jan-05	SMC2	Downstream sample site Ramona Grasslands	STREAM_PARAM	Stream Width	14		FT
24-Jan-05	SMC2	Downstream sample site Ramona Grasslands	STREAM_PARAM	Stream Width	11		FT
14-Feb-05	SMC2	Downstream sample site Ramona Grasslands	STREAM_PARAM	Stream Width	22.4		FT

Sample Date	Station	Source Description	Analysis	Analyte name	Value	MDL*	Units
28-Feb-05	SMC2	Downstream sample site Ramona Grasslands	STREAM_PARAM	Stream Width	19		FT
15-Mar-05	SMC2	Downstream sample site Ramona Grasslands	STREAM_PARAM	Stream Width	17		FT
28-Mar-05	SMC2	Downstream sample site Ramona Grasslands	STREAM_PARAM	Stream Width	13		FT
11-Apr-05	SMC2	Downstream sample site Ramona Grasslands	STREAM_PARAM	Stream Width	13		FT
25-Apr-05	SMC2	Downstream sample site Ramona Grasslands	STREAM_PARAM	Stream Width	10		FT
9-May-05	SMC2	Downstream sample site Ramona Grasslands	STREAM_PARAM	Stream Width	9		FT
23-May-05	SMC2	Downstream sample site Ramona Grasslands	STREAM_PARAM	Stream Width	1.9		FT
13-Mar-06	SMC2	Downstream sample site Ramona Grasslands	STREAM_PARAM	Stream Width	1.03		FT
3-Jan-05	SMC2	Downstream sample site Ramona Grasslands	ANIONS_IC	Sulfate	41.3	0.5	MG/L
28-Feb-05	SMC2	Downstream sample site Ramona Grasslands	ANIONS_IC	Sulfate	83.9	0.5	MG/L
11-Apr-05	SMC2	Downstream sample site Ramona Grasslands	ANIONS_IC	Sulfate	122	0.5	MG/L
25-Apr-05	SMC2	Downstream sample site Ramona Grasslands	ANIONS_IC	Sulfate	123	0.5	MG/L
9-May-05	SMC2	Downstream sample site Ramona Grasslands	ANIONS_IC	Sulfate	119	0.5	MG/L
23-May-05	SMC2	Downstream sample site Ramona Grasslands	ANIONS_IC	Sulfate	125	0.5	MG/L
13-Mar-06	SMC2	Downstream sample site Ramona Grasslands	ANIONS_IC	Sulfate	72.8	0.5	MG/L
3-Jan-05	SMC2	Downstream sample site Ramona Grasslands	STREAM_PARAM	Temperature degrees C	8.38		DEGREE_C
24-Jan-05	SMC2	Downstream sample site Ramona Grasslands	STREAM_PARAM	Temperature degrees C	10.6		DEGREE_C
14-Feb-05	SMC2	Downstream sample site Ramona Grasslands	STREAM_PARAM	Temperature degrees C	11.8		DEGREE_C
28-Feb-05	SMC2	Downstream sample site Ramona Grasslands	STREAM_PARAM	Temperature degrees C	11.4		DEGREE_C

Sample Date	Station	Source Description	Analysis	Analyte name	Value	MDL*	Units
15-Mar-05	SMC2	Downstream sample site Ramona Grasslands	STREAM_PARAM	Temperature degrees C	10.5		DEGREE_C
28-Mar-05	SMC2	Downstream sample site Ramona Grasslands	STREAM_PARAM	Temperature degrees C	13		DEGREE_C
11-Apr-05	SMC2	Downstream sample site Ramona Grasslands	STREAM_PARAM	Temperature degrees C	10.9		DEGREE_C
25-Apr-05	SMC2	Downstream sample site Ramona Grasslands	STREAM_PARAM	Temperature degrees C	15.5		DEGREE_C
9-May-05	SMC2	Downstream sample site Ramona Grasslands	STREAM_PARAM	Temperature degrees C	16.2		DEGREE_C
23-May-05	SMC2	Downstream sample site Ramona Grasslands	STREAM_PARAM	Temperature degrees C	21.3		DEGREE_C
13-Mar-06	SMC2	Downstream sample site Ramona Grasslands	STREAM_PARAM	Temperature degrees C	8.9		DEGREE_C
3-Jan-05	SMC2	Downstream sample site Ramona Grasslands	QT_CST	Total Coliform	9600		/100 ML
24-Jan-05	SMC2	Downstream sample site Ramona Grasslands	QT_CST	Total Coliform	4900		/100 ML
14-Feb-05	SMC2	Downstream sample site Ramona Grasslands	QT_CST	Total Coliform	55000		/100 ML
28-Feb-05	SMC2	Downstream sample site Ramona Grasslands	QT_CST	Total Coliform	15000		/100 ML
15-Mar-05	SMC2	Downstream sample site Ramona Grasslands	QT_CST	Total Coliform	18000		/100 ML
28-Mar-05	SMC2	Downstream sample site Ramona Grasslands	QT_CST	Total Coliform	2100		/100 ML
11-Apr-05	SMC2	Downstream sample site Ramona Grasslands	QT_CST	Total Coliform	8100		/100 ML
25-Apr-05	SMC2	Downstream sample site Ramona Grasslands	QT_CST	Total Coliform	16000		/100 ML
9-May-05	SMC2	Downstream sample site Ramona Grasslands	QT_CST	Total Coliform	10000		/100 ML
23-May-05	SMC2	Downstream sample site Ramona Grasslands	QT_CST	Total Coliform	26000		/100 ML
13-Mar-06	SMC2	Downstream sample site Ramona Grasslands	QT_CST	Total Coliform	26000		/100 ML
3-Jan-05	SMC2	Downstream sample site Ramona Grasslands	SOLIDS_TDS	Total Dissolved Solids	421	10	MG/L

Sample Date	Station	Source Description	Analysis	Analyte name	Value	MDL*	Units
24-Jan-05	SMC2	Downstream sample site Ramona Grasslands	SOLIDS_TDS	Total Dissolved Solids	793	10	MG/L
14-Feb-05	SMC2	Downstream sample site Ramona Grasslands	SOLIDS_TDS	Total Dissolved Solids	599	10	MG/L
28-Feb-05	SMC2	Downstream sample site Ramona Grasslands	SOLIDS_TDS	Total Dissolved Solids	613	10	MG/L
15-Mar-05	SMC2	Downstream sample site Ramona Grasslands	SOLIDS_TDS	Total Dissolved Solids	677	10	MG/L
28-Mar-05	SMC2	Downstream sample site Ramona Grasslands	SOLIDS_TDS	Total Dissolved Solids	41	10	MG/L
11-Apr-05	SMC2	Downstream sample site Ramona Grasslands	SOLIDS_TDS	Total Dissolved Solids	769	10	MG/L
25-Apr-05	SMC2	Downstream sample site Ramona Grasslands	SOLIDS_TDS	Total Dissolved Solids	785	10	MG/L
9-May-05	SMC2	Downstream sample site Ramona Grasslands	SOLIDS_TDS	Total Dissolved Solids	765	10	MG/L
23-May-05	SMC2	Downstream sample site Ramona Grasslands	SOLIDS_TDS	Total Dissolved Solids	1420	10	MG/L
13-Mar-06	SMC2	Downstream sample site Ramona Grasslands	SOLIDS_TDS	Total Dissolved Solids	665	10	MG/L
3-Jan-05	SMC2	Downstream sample site Ramona Grasslands	NIT_PHOS_UV	Total Nitrogen	1.95	0.22	MG/L
24-Jan-05	SMC2	Downstream sample site Ramona Grasslands	NIT_PHOS_UV	Total Nitrogen	6.3	0.22	MG/L
14-Feb-05	SMC2	Downstream sample site Ramona Grasslands	NIT_PHOS_UV	Total Nitrogen	5.69	0.22	MG/L
28-Feb-05	SMC2	Downstream sample site Ramona Grasslands	NIT_PHOS_UV	Total Nitrogen	8.83	0.22	MG/L
15-Mar-05	SMC2	Downstream sample site Ramona Grasslands	NIT_PHOS_UV	Total Nitrogen	8.91	0.22	MG/L
28-Mar-05	SMC2	Downstream sample site Ramona Grasslands	NIT_PHOS_UV	Total Nitrogen	7.66	0.22	MG/L
11-Apr-05	SMC2	Downstream sample site Ramona Grasslands	NIT_PHOS_UV	Total Nitrogen	8.44	0.22	MG/L
25-Apr-05	SMC2	Downstream sample site Ramona Grasslands	NIT_PHOS_UV	Total Nitrogen	7.85	0.22	MG/L
9-May-05	SMC2	Downstream sample site Ramona Grasslands	NIT_PHOS_UV	Total Nitrogen	6.62	0.22	MG/L

Sample Date	Station	Source Description	Analysis	Analyte name	Value	MDL*	Units
23-May-05	SMC2	Downstream sample site Ramona Grasslands	NIT_PHOS_UV	Total Nitrogen		0.22	MG/L
13-Mar-06	SMC2	Downstream sample site Ramona Grasslands	NIT_PHOS_UV	Total Nitrogen	1.37	0.156	MG/L
3-Jan-05	SMC2	Downstream sample site Ramona Grasslands	SOLIDS_TSS	Total Suspended Solids (TSS)	11.8		MG/L
24-Jan-05	SMC2	Downstream sample site Ramona Grasslands	SOLIDS_TSS	Total Suspended Solids (TSS)	1.55	1	MG/L
14-Feb-05	SMC2	Downstream sample site Ramona Grasslands	SOLIDS_TSS	Total Suspended Solids (TSS)	20	2	MG/L
28-Feb-05	SMC2	Downstream sample site Ramona Grasslands	SOLIDS_TSS	Total Suspended Solids (TSS)	58		MG/L
15-Mar-05	SMC2	Downstream sample site Ramona Grasslands	SOLIDS_TSS	Total Suspended Solids (TSS)	97.1	1	MG/L
28-Mar-05	SMC2	Downstream sample site Ramona Grasslands	SOLIDS_TSS	Total Suspended Solids (TSS)	15.1		MG/L
11-Apr-05	SMC2	Downstream sample site Ramona Grasslands	SOLIDS_TSS	Total Suspended Solids (TSS)	2		MG/L
25-Apr-05	SMC2	Downstream sample site Ramona Grasslands	SOLIDS_TSS	Total Suspended Solids (TSS)	4.5		MG/L
9-May-05	SMC2	Downstream sample site Ramona Grasslands	SOLIDS_TSS	Total Suspended Solids (TSS)	2.1		MG/L
23-May-05	SMC2	Downstream sample site Ramona Grasslands	SOLIDS_TSS	Total Suspended Solids (TSS)	21.9		MG/L
13-Mar-06	SMC2	Downstream sample site Ramona Grasslands	SOLIDS_TSS	Total Suspended Solids (TSS)	29.6		MG/L
3-Jan-05	SMC3	Santa Maria Creek Station 3 Mid Point	NH3_NO3_NO2_AUT	Ammonia-N	0.221	0.04	MG/L
24-Jan-05	SMC3	Santa Maria Creek Station 3 Mid Point	NH3_NO3_NO2_AUT	Ammonia-N	0.191	0.04	MG/L
14-Feb-05	SMC3	Santa Maria Creek Station 3 Mid Point	NH3_NO3_NO2_AUT	Ammonia-N	0.912	0.04	MG/L
28-Feb-05	SMC3	Santa Maria Creek Station 3 Mid Point	NH3_NO3_NO2_AUT	Ammonia-N	0.678	0.04	MG/L
15-Mar-05	SMC3	Santa Maria Creek Station 3 Mid Point	NH3_NO3_NO2_AUT	Ammonia-N	0.07	0.04	MG/L
28-Mar-05	SMC3	Santa Maria Creek Station 3 Mid Point	NH3_NO3_NO2_AUT	Ammonia-N	0.771	0.04	MG/L

Sample Date	Station	Source Description	Analysis	Analyte name	Value	MDL*	Units
11-Apr-05	SMC3	Santa Maria Creek Station 3 Mid Point	NH3_NO3_NO2_AUT	Ammonia-N	ND	0.04	MG/L
25-Apr-05	SMC3	Santa Maria Creek Station 3 Mid Point	NH3_NO3_NO2_AUT	Ammonia-N	0.051	0.04	MG/L
9-May-05	SMC3	Santa Maria Creek Station 3 Mid Point	NH3_NO3_NO2_AUT	Ammonia-N	ND	0.04	MG/L
23-May-05	SMC3	Santa Maria Creek Station 3 Mid Point	NH3_NO3_NO2_AUT	Ammonia-N	ND	0.04	MG/L
13-Mar-06	SMC3	Santa Maria Creek Station 3 Mid Point	NH3_NO3_NO2_AUT	Ammonia-N	0.059	0.031	MG/L
3-Jan-05	SMC3	Santa Maria Creek Station 3 Mid Point	ANIONS_IC	Bromide	0.167	0.1	MG/L
28-Feb-05	SMC3	Santa Maria Creek Station 3 Mid Point	ANIONS_IC	Bromide	0.25	0.1	MG/L
11-Apr-05	SMC3	Santa Maria Creek Station 3 Mid Point	ANIONS_IC	Bromide	0.393	0.1	MG/L
25-Apr-05	SMC3	Santa Maria Creek Station 3 Mid Point	ANIONS_IC	Bromide	0.381	0.1	MG/L
9-May-05	SMC3	Santa Maria Creek Station 3 Mid Point	ANIONS_IC	Bromide	0.401	0.1	MG/L
23-May-05	SMC3	Santa Maria Creek Station 3 Mid Point	ANIONS_IC	Bromide	0.489	0.1	MG/L
13-Mar-06	SMC3	Santa Maria Creek Station 3 Mid Point	ANIONS_IC	Bromide	0.11	0.1	MG/L
3-Jan-05	SMC3	Santa Maria Creek Station 3 Mid Point	ANIONS_IC	Chloride	75.9	0.5	MG/L
28-Feb-05	SMC3	Santa Maria Creek Station 3 Mid Point	ANIONS_IC	Chloride	105	1	MG/L
11-Apr-05	SMC3	Santa Maria Creek Station 3 Mid Point	ANIONS_IC	Chloride	157	1	MG/L
25-Apr-05	SMC3	Santa Maria Creek Station 3 Mid Point	ANIONS_IC	Chloride	158	1	MG/L
9-May-05	SMC3	Santa Maria Creek Station 3 Mid Point	ANIONS_IC	Chloride	153	1	MG/L
23-May-05	SMC3	Santa Maria Creek Station 3 Mid Point	ANIONS_IC	Chloride	131	1	MG/L
13-Mar-06	SMC3	Santa Maria Creek Station 3 Mid Point	ANIONS_IC	Chloride	57	0.5	MG/L

Sample Date	Station	Source Description	Analysis	Analyte name	Value	MDL*	Units
3-Jan-05	SMC3	Santa Maria Creek Station 3 Mid Point	STREAM_PARAM	Conductivity	677		US/CM
24-Jan-05	SMC3	Santa Maria Creek Station 3 Mid Point	STREAM_PARAM	Conductivity	1400		US/CM
14-Feb-05	SMC3	Santa Maria Creek Station 3 Mid Point	STREAM_PARAM	Conductivity	965		US/CM
28-Feb-05	SMC3	Santa Maria Creek Station 3 Mid Point	STREAM_PARAM	Conductivity	922		US/CM
15-Mar-05	SMC3	Santa Maria Creek Station 3 Mid Point	STREAM_PARAM	Conductivity	1180		US/CM
28-Mar-05	SMC3	Santa Maria Creek Station 3 Mid Point	STREAM_PARAM	Conductivity	1200		US/CM
11-Apr-05	SMC3	Santa Maria Creek Station 3 Mid Point	STREAM_PARAM	Conductivity	1260		US/CM
25-Apr-05	SMC3	Santa Maria Creek Station 3 Mid Point	STREAM_PARAM	Conductivity	1040		US/CM
9-May-05	SMC3	Santa Maria Creek Station 3 Mid Point	STREAM_PARAM	Conductivity	1220		US/CM
23-May-05	SMC3	Santa Maria Creek Station 3 Mid Point	STREAM_PARAM	Conductivity	915		US/CM
13-Mar-06	SMC3	Santa Maria Creek Station 3 Mid Point	STREAM_PARAM	Conductivity	439		US/CM
3-Jan-05	SMC3	Santa Maria Creek Station 3 Mid Point	STREAM_PARAM	Dissolved Oxygen	9.98		MG/L
24-Jan-05	SMC3	Santa Maria Creek Station 3 Mid Point	STREAM_PARAM	Dissolved Oxygen	9.21		MG/L
14-Feb-05	SMC3	Santa Maria Creek Station 3 Mid Point	STREAM_PARAM	Dissolved Oxygen	8.45		MG/L
28-Feb-05	SMC3	Santa Maria Creek Station 3 Mid Point	STREAM_PARAM	Dissolved Oxygen	9.23		MG/L
15-Mar-05	SMC3	Santa Maria Creek Station 3 Mid Point	STREAM_PARAM	Dissolved Oxygen	10.3		MG/L
28-Mar-05	SMC3	Santa Maria Creek Station 3 Mid Point	STREAM_PARAM	Dissolved Oxygen	9.66		MG/L
11-Apr-05	SMC3	Santa Maria Creek Station 3 Mid Point	STREAM_PARAM	Dissolved Oxygen	10.4		MG/L
25-Apr-05	SMC3	Santa Maria Creek Station 3 Mid Point	STREAM_PARAM	Dissolved Oxygen	9.64		MG/L

Sample Date	Station	Source Description	Analysis	Analyte name	Value	MDL*	Units
9-May-05	SMC3	Santa Maria Creek Station 3 Mid Point	STREAM_PARAM	Dissolved Oxygen	8.16		MG/L
23-May-05	SMC3	Santa Maria Creek Station 3 Mid Point	STREAM_PARAM	Dissolved Oxygen	5.44		MG/L
13-Mar-06	SMC3	Santa Maria Creek Station 3 Mid Point	STREAM_PARAM	Dissolved Oxygen	9.8		MG/L
3-Jan-05	SMC3	Santa Maria Creek Station 3 Mid Point	QT_CST	Enterococcus	70		/100 ML
24-Jan-05	SMC3	Santa Maria Creek Station 3 Mid Point	QT_CST	Enterococcus	27		/100 ML
14-Feb-05	SMC3	Santa Maria Creek Station 3 Mid Point	QT_CST	Enterococcus	1100		/100 ML
28-Feb-05	SMC3	Santa Maria Creek Station 3 Mid Point	QT_CST	Enterococcus	82		/100 ML
15-Mar-05	SMC3	Santa Maria Creek Station 3 Mid Point	QT_CST	Enterococcus	130		/100 ML
28-Mar-05	SMC3	Santa Maria Creek Station 3 Mid Point	QT_CST	Enterococcus	110		/100 ML
11-Apr-05	SMC3	Santa Maria Creek Station 3 Mid Point	QT_CST	Enterococcus	15		/100 ML
25-Apr-05	SMC3	Santa Maria Creek Station 3 Mid Point	QT_CST	Enterococcus	84		/100 ML
9-May-05	SMC3	Santa Maria Creek Station 3 Mid Point	QT_CST	Enterococcus	93		/100 ML
23-May-05	SMC3	Santa Maria Creek Station 3 Mid Point	QT_CST	Enterococcus	520		/100 ML
13-Mar-06	SMC3	Santa Maria Creek Station 3 Mid Point	QT_CST	Enterococcus	980		/100 ML
3-Jan-05	SMC3	Santa Maria Creek Station 3 Mid Point	STREAM_PARAM	Equipment Number	39300		.
24-Jan-05	SMC3	Santa Maria Creek Station 3 Mid Point	STREAM_PARAM	Equipment Number	39300		.
14-Feb-05	SMC3	Santa Maria Creek Station 3 Mid Point	STREAM_PARAM	Equipment Number	39300		.
28-Feb-05	SMC3	Santa Maria Creek Station 3 Mid Point	STREAM_PARAM	Equipment Number	39300		.
15-Mar-05	SMC3	Santa Maria Creek Station 3 Mid Point	STREAM_PARAM	Equipment Number	39300		.

Sample Date	Station	Source Description	Analysis	Analyte name	Value	MDL*	Units
28-Mar-05	SMC3	Santa Maria Creek Station 3 Mid Point	STREAM_PARAM	Equipment Number	39300		.
11-Apr-05	SMC3	Santa Maria Creek Station 3 Mid Point	STREAM_PARAM	Equipment Number	39300		.
25-Apr-05	SMC3	Santa Maria Creek Station 3 Mid Point	STREAM_PARAM	Equipment Number	39300		.
9-May-05	SMC3	Santa Maria Creek Station 3 Mid Point	STREAM_PARAM	Equipment Number	39300		.
23-May-05	SMC3	Santa Maria Creek Station 3 Mid Point	STREAM_PARAM	Equipment Number	39300		.
3-Jan-05	SMC3	Santa Maria Creek Station 3 Mid Point	QT_CST	Escherichia Coli	630		/100 ML
24-Jan-05	SMC3	Santa Maria Creek Station 3 Mid Point	QT_CST	Escherichia Coli	410		/100 ML
14-Feb-05	SMC3	Santa Maria Creek Station 3 Mid Point	QT_CST	Escherichia Coli	4700		/100 ML
28-Feb-05	SMC3	Santa Maria Creek Station 3 Mid Point	QT_CST	Escherichia Coli	720		/100 ML
15-Mar-05	SMC3	Santa Maria Creek Station 3 Mid Point	QT_CST	Escherichia Coli	740		/100 ML
28-Mar-05	SMC3	Santa Maria Creek Station 3 Mid Point	QT_CST	Escherichia Coli	630		/100 ML
11-Apr-05	SMC3	Santa Maria Creek Station 3 Mid Point	QT_CST	Escherichia Coli	520		/100 ML
25-Apr-05	SMC3	Santa Maria Creek Station 3 Mid Point	QT_CST	Escherichia Coli	630		/100 ML
9-May-05	SMC3	Santa Maria Creek Station 3 Mid Point	QT_CST	Escherichia Coli	200		/100 ML
23-May-05	SMC3	Santa Maria Creek Station 3 Mid Point	QT_CST	Escherichia Coli	< 100		/100 ML
13-Mar-06	SMC3	Santa Maria Creek Station 3 Mid Point	QT_CST	Escherichia Coli	1900		/100 ML
3-Jan-05	SMC3	Santa Maria Creek Station 3 Mid Point	ANIONS_IC	Nitrate	4.85	0.2	MG/L
24-Jan-05	SMC3	Santa Maria Creek Station 3 Mid Point	NH3_NO3_NO2_AUT	Nitrate	24.1	0.6	MG/L
14-Feb-05	SMC3	Santa Maria Creek Station 3 Mid Point	NH3_NO3_NO2_AUT	Nitrate	12.3	0.6	MG/L

Sample Date	Station	Source Description	Analysis	Analyte name	Value	MDL*	Units
28-Feb-05	SMC3	Santa Maria Creek Station 3 Mid Point	NH3_NO3_NO2_AUT	Nitrate	25.8	0.6	MG/L
15-Mar-05	SMC3	Santa Maria Creek Station 3 Mid Point	NH3_NO3_NO2_AUT	Nitrate	34.4	1.2	MG/L
28-Mar-05	SMC3	Santa Maria Creek Station 3 Mid Point	NH3_NO3_NO2_AUT	Nitrate	30.4	0.6	MG/L
11-Apr-05	SMC3	Santa Maria Creek Station 3 Mid Point	ANIONS_IC	Nitrate	36.1	0.6	MG/L
25-Apr-05	SMC3	Santa Maria Creek Station 3 Mid Point	NH3_NO3_NO2_AUT	Nitrate	> 25	0.3	MG/L
9-May-05	SMC3	Santa Maria Creek Station 3 Mid Point	ANIONS_IC	Nitrate	29.2	0.2	MG/L
9-May-05	SMC3	Santa Maria Creek Station 3 Mid Point	NH3_NO3_NO2_AUT	Nitrate	26.9	0.6	MG/L
23-May-05	SMC3	Santa Maria Creek Station 3 Mid Point	NH3_NO3_NO2_AUT	Nitrate	0.146	0.06	MG/L
13-Mar-06	SMC3	Santa Maria Creek Station 3 Mid Point	ANIONS_IC	Nitrate	5.31	0.4	MG/L
13-Mar-06	SMC3	Santa Maria Creek Station 3 Mid Point	NH3_NO3_NO2_AUT	Nitrate	4.76	0.078	MG/L
3-Jan-05	SMC3	Santa Maria Creek Station 3 Mid Point	NH3_NO3_NO2_AUT	Nitrate & Nitrite	4.6	0.06	MG/L
24-Jan-05	SMC3	Santa Maria Creek Station 3 Mid Point	NH3_NO3_NO2_AUT	Nitrate & Nitrite	24.5	0.6	MG/L
14-Feb-05	SMC3	Santa Maria Creek Station 3 Mid Point	NH3_NO3_NO2_AUT	Nitrate & Nitrite	12.7	0.6	MG/L
28-Feb-05	SMC3	Santa Maria Creek Station 3 Mid Point	NH3_NO3_NO2_AUT	Nitrate & Nitrite	26.8	0.6	MG/L
15-Mar-05	SMC3	Santa Maria Creek Station 3 Mid Point	NH3_NO3_NO2_AUT	Nitrate & Nitrite	35	1.2	MG/L
28-Mar-05	SMC3	Santa Maria Creek Station 3 Mid Point	NH3_NO3_NO2_AUT	Nitrate & Nitrite	30.9	0.6	MG/L
25-Apr-05	SMC3	Santa Maria Creek Station 3 Mid Point	NH3_NO3_NO2_AUT	Nitrate & Nitrite	> 25	0.3	MG/L
9-May-05	SMC3	Santa Maria Creek Station 3 Mid Point	NH3_NO3_NO2_AUT	Nitrate & Nitrite	27.1	0.6	MG/L
23-May-05	SMC3	Santa Maria Creek Station 3 Mid Point	NH3_NO3_NO2_AUT	Nitrate & Nitrite	0.185	0.06	MG/L

Sample Date	Station	Source Description	Analysis	Analyte name	Value	MDL*	Units
13-Mar-06	SMC3	Santa Maria Creek Station 3 Mid Point	NH3_NO3_NO2_AUT	Nitrate & Nitrite	4.88	0.078	MG/L
3-Jan-05	SMC3	Santa Maria Creek Station 3 Mid Point	NH3_NO3_NO2_AUT	Nitrite (NO2)	0.114	0.01	MG/L
24-Jan-05	SMC3	Santa Maria Creek Station 3 Mid Point	NH3_NO3_NO2_AUT	Nitrite (NO2)	0.383	0.01	MG/L
14-Feb-05	SMC3	Santa Maria Creek Station 3 Mid Point	NH3_NO3_NO2_AUT	Nitrite (NO2)	0.445	0.01	MG/L
28-Feb-05	SMC3	Santa Maria Creek Station 3 Mid Point	NH3_NO3_NO2_AUT	Nitrite (NO2)	1.09	0.05	MG/L
15-Mar-05	SMC3	Santa Maria Creek Station 3 Mid Point	NH3_NO3_NO2_AUT	Nitrite (NO2)	0.5	0.01	MG/L
28-Mar-05	SMC3	Santa Maria Creek Station 3 Mid Point	NH3_NO3_NO2_AUT	Nitrite (NO2)	0.392	0.01	MG/L
25-Apr-05	SMC3	Santa Maria Creek Station 3 Mid Point	NH3_NO3_NO2_AUT	Nitrite (NO2)	0.213	0.01	MG/L
9-May-05	SMC3	Santa Maria Creek Station 3 Mid Point	NH3_NO3_NO2_AUT	Nitrite (NO2)	0.184	0.01	MG/L
23-May-05	SMC3	Santa Maria Creek Station 3 Mid Point	NH3_NO3_NO2_AUT	Nitrite (NO2)	0.039	0.01	MG/L
13-Mar-06	SMC3	Santa Maria Creek Station 3 Mid Point	NH3_NO3_NO2_AUT	Nitrite (NO2)	0.119	0.016	MG/L
3-Jan-05	SMC3	Santa Maria Creek Station 3 Mid Point	ANIONS_IC	Ortho phosphates	1.45	0.2	MG/L
24-Jan-05	SMC3	Santa Maria Creek Station 3 Mid Point	ANIONS_IC_LOW	Ortho phosphates	1.64	0.2	MG/L
14-Feb-05	SMC3	Santa Maria Creek Station 3 Mid Point	ANIONS_IC_LOW	Ortho phosphates	3.51	0.05	MG/L
28-Feb-05	SMC3	Santa Maria Creek Station 3 Mid Point	ANIONS_IC_LOW	Ortho phosphates	2.65	0.15	MG/L
15-Mar-05	SMC3	Santa Maria Creek Station 3 Mid Point	ANIONS_IC_LOW	Ortho phosphates	1.55	0.05	MG/L
28-Mar-05	SMC3	Santa Maria Creek Station 3 Mid Point	ANIONS_IC_LOW	Ortho phosphates	1.89	0.05	MG/L
11-Apr-05	SMC3	Santa Maria Creek Station 3 Mid Point	ANIONS_IC	Ortho phosphates	1.3	0.2	MG/L
25-Apr-05	SMC3	Santa Maria Creek Station 3 Mid Point	ANIONS_IC_LOW	Ortho phosphates	1.34	0.1	MG/L

Sample Date	Station	Source Description	Analysis	Analyte name	Value	MDL*	Units
9-May-05	SMC3	Santa Maria Creek Station 3 Mid Point	ANIONS_IC	Ortho phosphates	1.22	0.2	MG/L
23-May-05	SMC3	Santa Maria Creek Station 3 Mid Point	ANIONS_IC	Ortho phosphates	1.01	0.2	MG/L
23-May-05	SMC3	Santa Maria Creek Station 3 Mid Point	ANIONS_IC_LOW	Ortho phosphates	1.16	0.1	MG/L
13-Mar-06	SMC3	Santa Maria Creek Station 3 Mid Point	ANIONS_IC	Ortho phosphates	1.22	0.2	MG/L
3-Jan-05	SMC3	Santa Maria Creek Station 3 Mid Point	STREAM_PARAM	Oxidation- Reduction Potential	384		MV
24-Jan-05	SMC3	Santa Maria Creek Station 3 Mid Point	STREAM_PARAM	Oxidation- Reduction Potential	378		MV
14-Feb-05	SMC3	Santa Maria Creek Station 3 Mid Point	STREAM_PARAM	Oxidation- Reduction Potential	348		MV
28-Feb-05	SMC3	Santa Maria Creek Station 3 Mid Point	STREAM_PARAM	Oxidation- Reduction Potential	348		MV
15-Mar-05	SMC3	Santa Maria Creek Station 3 Mid Point	STREAM_PARAM	Oxidation- Reduction Potential	388		MV
28-Mar-05	SMC3	Santa Maria Creek Station 3 Mid Point	STREAM_PARAM	Oxidation- Reduction Potential	419		MV
11-Apr-05	SMC3	Santa Maria Creek Station 3 Mid Point	STREAM_PARAM	Oxidation- Reduction Potential	336		MV
25-Apr-05	SMC3	Santa Maria Creek Station 3 Mid Point	STREAM_PARAM	Oxidation- Reduction Potential	393		MV
9-May-05	SMC3	Santa Maria Creek Station 3 Mid Point	STREAM_PARAM	Oxidation- Reduction Potential	303		MV
23-May-05	SMC3	Santa Maria Creek Station 3 Mid Point	STREAM_PARAM	Oxidation- Reduction Potential	176		MV

Sample Date	Station	Source Description	Analysis	Analyte name	Value	MDL*	Units
13-Mar-06	SMC3	Santa Maria Creek Station 3 Mid Point	STREAM_PARAM	Oxidation- Reduction Potential	389		MV
3-Jan-05	SMC3	Santa Maria Creek Station 3 Mid Point	STREAM_PARAM	pH	7.67		PH
24-Jan-05	SMC3	Santa Maria Creek Station 3 Mid Point	STREAM_PARAM	pH	7.85		PH
14-Feb-05	SMC3	Santa Maria Creek Station 3 Mid Point	STREAM_PARAM	pH	7.63		PH
28-Feb-05	SMC3	Santa Maria Creek Station 3 Mid Point	STREAM_PARAM	pH	7.75		PH
15-Mar-05	SMC3	Santa Maria Creek Station 3 Mid Point	STREAM_PARAM	pH	7.91		PH
28-Mar-05	SMC3	Santa Maria Creek Station 3 Mid Point	STREAM_PARAM	pH	8		PH
11-Apr-05	SMC3	Santa Maria Creek Station 3 Mid Point	STREAM_PARAM	pH	7.99		PH
25-Apr-05	SMC3	Santa Maria Creek Station 3 Mid Point	STREAM_PARAM	pH	8.33		PH
9-May-05	SMC3	Santa Maria Creek Station 3 Mid Point	STREAM_PARAM	pH	8.35		PH
23-May-05	SMC3	Santa Maria Creek Station 3 Mid Point	STREAM_PARAM	pH	7.75		PH
13-Mar-06	SMC3	Santa Maria Creek Station 3 Mid Point	STREAM_PARAM	pH	7.71		PH
3-Jan-05	SMC3	Santa Maria Creek Station 3 Mid Point	NIT_PHOS_UV	Phosphorus	0.594	0.084	MG/L
24-Jan-05	SMC3	Santa Maria Creek Station 3 Mid Point	NIT_PHOS_UV	Phosphorus	0.685	0.084	MG/L
14-Feb-05	SMC3	Santa Maria Creek Station 3 Mid Point	NIT_PHOS_UV	Phosphorus	1.32	0.084	MG/L
28-Feb-05	SMC3	Santa Maria Creek Station 3 Mid Point	NIT_PHOS_UV	Phosphorus	0.992	0.084	MG/L
15-Mar-05	SMC3	Santa Maria Creek Station 3 Mid Point	NIT_PHOS_UV	Phosphorus	0.834	0.084	MG/L
28-Mar-05	SMC3	Santa Maria Creek Station 3 Mid Point	NIT_PHOS_UV	Phosphorus	0.793	0.084	MG/L
11-Apr-05	SMC3	Santa Maria Creek Station 3 Mid Point	NIT_PHOS_UV	Phosphorus	0.612	0.084	MG/L

Sample Date	Station	Source Description	Analysis	Analyte name	Value	MDL*	Units
25-Apr-05	SMC3	Santa Maria Creek Station 3 Mid Point	NIT_PHOS_UV	Phosphorus	0.538	0.084	MG/L
9-May-05	SMC3	Santa Maria Creek Station 3 Mid Point	NIT_PHOS_UV	Phosphorus	0.585	0.084	MG/L
23-May-05	SMC3	Santa Maria Creek Station 3 Mid Point	NIT_PHOS_UV	Phosphorus		0.084	MG/L
13-Mar-06	SMC3	Santa Maria Creek Station 3 Mid Point	NIT_PHOS_UV	Phosphorus	0.497	0.078	MG/L
3-Jan-05	SMC3	Santa Maria Creek Station 3 Mid Point	STREAM_PARAM	Stream Average Depth	0.33		FT
24-Jan-05	SMC3	Santa Maria Creek Station 3 Mid Point	STREAM_PARAM	Stream Average Depth	0.34		FT
14-Feb-05	SMC3	Santa Maria Creek Station 3 Mid Point	STREAM_PARAM	Stream Average Depth	1.08		FT
28-Feb-05	SMC3	Santa Maria Creek Station 3 Mid Point	STREAM_PARAM	Stream Average Depth	0.97		FT
15-Mar-05	SMC3	Santa Maria Creek Station 3 Mid Point	STREAM_PARAM	Stream Average Depth	0.77		FT
28-Mar-05	SMC3	Santa Maria Creek Station 3 Mid Point	STREAM_PARAM	Stream Average Depth	0.59		FT
11-Apr-05	SMC3	Santa Maria Creek Station 3 Mid Point	STREAM_PARAM	Stream Average Depth	0.29		FT
25-Apr-05	SMC3	Santa Maria Creek Station 3 Mid Point	STREAM_PARAM	Stream Average Depth	0.26		FT
9-May-05	SMC3	Santa Maria Creek Station 3 Mid Point	STREAM_PARAM	Stream Average Depth	0.3		FT
23-May-05	SMC3	Santa Maria Creek Station 3 Mid Point	STREAM_PARAM	Stream Average Depth	0.1		FT
13-Mar-06	SMC3	Santa Maria Creek Station 3 Mid Point	STREAM_PARAM	Stream Average Depth	0.23		FT
3-Jan-05	SMC3	Santa Maria Creek Station 3 Mid Point	STREAM_PARAM	Stream Flow	3.16		FT3/S
24-Jan-05	SMC3	Santa Maria Creek Station 3 Mid Point	STREAM_PARAM	Stream Flow	3.1		FT3/S
14-Feb-05	SMC3	Santa Maria Creek Station 3 Mid Point	STREAM_PARAM	Stream Flow	25.4		FT3/S
28-Feb-05	SMC3	Santa Maria Creek Station 3 Mid Point	STREAM_PARAM	Stream Flow	33.6		FT3/S

Sample Date	Station	Source Description	Analysis	Analyte name	Value	MDL*	Units
15-Mar-05	SMC3	Santa Maria Creek Station 3 Mid Point	STREAM_PARAM	Stream Flow	12.5		FT3/S
28-Mar-05	SMC3	Santa Maria Creek Station 3 Mid Point	STREAM_PARAM	Stream Flow	11.9		FT3/S
11-Apr-05	SMC3	Santa Maria Creek Station 3 Mid Point	STREAM_PARAM	Stream Flow	3.8		FT3/S
25-Apr-05	SMC3	Santa Maria Creek Station 3 Mid Point	STREAM_PARAM	Stream Flow	3.38		FT3/S
9-May-05	SMC3	Santa Maria Creek Station 3 Mid Point	STREAM_PARAM	Stream Flow	3.7		FT3/S
23-May-05	SMC3	Santa Maria Creek Station 3 Mid Point	STREAM_PARAM	Stream Flow	0.12		FT3/S
13-Mar-06	SMC3	Santa Maria Creek Station 3 Mid Point	STREAM_PARAM	Stream Flow	0.35		FT3/S
3-Jan-05	SMC3	Santa Maria Creek Station 3 Mid Point	STREAM_PARAM	Stream Width	5.8		FT
24-Jan-05	SMC3	Santa Maria Creek Station 3 Mid Point	STREAM_PARAM	Stream Width	8		FT
14-Feb-05	SMC3	Santa Maria Creek Station 3 Mid Point	STREAM_PARAM	Stream Width	14.5		FT
28-Feb-05	SMC3	Santa Maria Creek Station 3 Mid Point	STREAM_PARAM	Stream Width	16		FT
15-Mar-05	SMC3	Santa Maria Creek Station 3 Mid Point	STREAM_PARAM	Stream Width	10		FT
28-Mar-05	SMC3	Santa Maria Creek Station 3 Mid Point	STREAM_PARAM	Stream Width	11		FT
11-Apr-05	SMC3	Santa Maria Creek Station 3 Mid Point	STREAM_PARAM	Stream Width	11		FT
25-Apr-05	SMC3	Santa Maria Creek Station 3 Mid Point	STREAM_PARAM	Stream Width	9.4		FT
9-May-05	SMC3	Santa Maria Creek Station 3 Mid Point	STREAM_PARAM	Stream Width	9		FT
23-May-05	SMC3	Santa Maria Creek Station 3 Mid Point	STREAM_PARAM	Stream Width	1.3		FT
13-Mar-06	SMC3	Santa Maria Creek Station 3 Mid Point	STREAM_PARAM	Stream Width	0.8		FT
3-Jan-05	SMC3	Santa Maria Creek Station 3 Mid Point	ANIONS_IC	Sulfate	43.8	0.5	MG/L

Sample Date	Station	Source Description	Analysis	Analyte name	Value	MDL*	Units
28-Feb-05	SMC3	Santa Maria Creek Station 3 Mid Point	ANIONS_IC	Sulfate	83.4	0.5	MG/L
11-Apr-05	SMC3	Santa Maria Creek Station 3 Mid Point	ANIONS_IC	Sulfate	122	0.5	MG/L
25-Apr-05	SMC3	Santa Maria Creek Station 3 Mid Point	ANIONS_IC	Sulfate	124	0.5	MG/L
9-May-05	SMC3	Santa Maria Creek Station 3 Mid Point	ANIONS_IC	Sulfate	120	0.5	MG/L
23-May-05	SMC3	Santa Maria Creek Station 3 Mid Point	ANIONS_IC	Sulfate	53.8	0.5	MG/L
13-Mar-06	SMC3	Santa Maria Creek Station 3 Mid Point	ANIONS_IC	Sulfate	40.4	0.5	MG/L
3-Jan-05	SMC3	Santa Maria Creek Station 3 Mid Point	STREAM_PARAM	Temperature degrees C	8.12		DEGREE_C
24-Jan-05	SMC3	Santa Maria Creek Station 3 Mid Point	STREAM_PARAM	Temperature degrees C	11.1		DEGREE_C
14-Feb-05	SMC3	Santa Maria Creek Station 3 Mid Point	STREAM_PARAM	Temperature degrees C	11.7		DEGREE_C
28-Feb-05	SMC3	Santa Maria Creek Station 3 Mid Point	STREAM_PARAM	Temperature degrees C	11.5		DEGREE_C
15-Mar-05	SMC3	Santa Maria Creek Station 3 Mid Point	STREAM_PARAM	Temperature degrees C	10.1		DEGREE_C
28-Mar-05	SMC3	Santa Maria Creek Station 3 Mid Point	STREAM_PARAM	Temperature degrees C	12.5		DEGREE_C
11-Apr-05	SMC3	Santa Maria Creek Station 3 Mid Point	STREAM_PARAM	Temperature degrees C	10.2		DEGREE_C
25-Apr-05	SMC3	Santa Maria Creek Station 3 Mid Point	STREAM_PARAM	Temperature degrees C	14.8		DEGREE_C
9-May-05	SMC3	Santa Maria Creek Station 3 Mid Point	STREAM_PARAM	Temperature degrees C	17.8		DEGREE_C
23-May-05	SMC3	Santa Maria Creek Station 3 Mid Point	STREAM_PARAM	Temperature degrees C	18		DEGREE_C
3-Jan-05	SMC3	Santa Maria Creek Station 3 Mid Point	QT_CST	Total Coliform	13000		/100 ML
24-Jan-05	SMC3	Santa Maria Creek Station 3 Mid Point	QT_CST	Total Coliform	7100		/100 ML
14-Feb-05	SMC3	Santa Maria Creek Station 3 Mid Point	QT_CST	Total Coliform	37000		/100 ML

Sample Date	Station	Source Description	Analysis	Analyte name	Value	MDL*	Units
28-Feb-05	SMC3	Santa Maria Creek Station 3 Mid Point	QT_CST	Total Coliform	18000		/100 ML
15-Mar-05	SMC3	Santa Maria Creek Station 3 Mid Point	QT_CST	Total Coliform	28000		/100 ML
28-Mar-05	SMC3	Santa Maria Creek Station 3 Mid Point	QT_CST	Total Coliform	9600		/100 ML
11-Apr-05	SMC3	Santa Maria Creek Station 3 Mid Point	QT_CST	Total Coliform	9900		/100 ML
25-Apr-05	SMC3	Santa Maria Creek Station 3 Mid Point	QT_CST	Total Coliform	1200		/100 ML
9-May-05	SMC3	Santa Maria Creek Station 3 Mid Point	QT_CST	Total Coliform	19000		/100 ML
23-May-05	SMC3	Santa Maria Creek Station 3 Mid Point	QT_CST	Total Coliform	46000		/100 ML
13-Mar-06	SMC3	Santa Maria Creek Station 3 Mid Point	QT_CST	Total Coliform	41000		/100 ML
3-Jan-05	SMC3	Santa Maria Creek Station 3 Mid Point	SOLIDS_TDS	Total Dissolved Solids	397	10	MG/L
24-Jan-05	SMC3	Santa Maria Creek Station 3 Mid Point	SOLIDS_TDS	Total Dissolved Solids	848	10	MG/L
14-Feb-05	SMC3	Santa Maria Creek Station 3 Mid Point	SOLIDS_TDS	Total Dissolved Solids	619	10	MG/L
28-Feb-05	SMC3	Santa Maria Creek Station 3 Mid Point	SOLIDS_TDS	Total Dissolved Solids	614	10	MG/L
15-Mar-05	SMC3	Santa Maria Creek Station 3 Mid Point	SOLIDS_TDS	Total Dissolved Solids	734	10	MG/L
28-Mar-05	SMC3	Santa Maria Creek Station 3 Mid Point	SOLIDS_TDS	Total Dissolved Solids	784	10	MG/L
11-Apr-05	SMC3	Santa Maria Creek Station 3 Mid Point	SOLIDS_TDS	Total Dissolved Solids	786	10	MG/L

\*MDL = Method Detection Limit